



**Coastal Zone and
Estuarine Studies
Division**

**Northwest Fisheries
Science Center**


**National Marine
Fisheries Service**

Seattle, Washington

***Benthic invertebrates
in soft-substrate,
shallow-water habitats
in Lower Granite Reservoir,
1994-95***

by
Suzan S. Pool and
Richard D. Ledgerwood

July 1997



**BENTHIC INVERTEBRATES IN SOFT-SUBSTRATE, SHALLOW-WATER
HABITATS IN LOWER GRANITE RESERVOIR, 1994-95**

by

Suzan S. Pool
and
Richard D. Ledgerwood

Funded by

U.S. Army Corps of Engineers
Walla Walla District
201 North 3rd
Walla Walla, Washington 99362-1876
(Contract E86940115)

and

Coastal Zone and Estuarine Studies Division
Northwest Fisheries Science Center
National Marine Fisheries Service
National Oceanic and Atmospheric Administration
2725 Montlake Boulevard East
Seattle, Washington 98112-2097

July 1997

EXECUTIVE SUMMARY

In an effort to recover dwindling Snake River salmon runs, a drawdown or lowering of the water level in Lower Granite Reservoir has been proposed. Some fisheries managers believe that reservoir drawdown would help restoration efforts for depleted upriver stocks of Snake River chinook (*Oncorhynchus tshawytscha*) and sockeye salmon (*O. nerka*). The drawdown would simulate a free-flowing river, which would theoretically decrease downstream travel time and thus increase the survival of juvenile salmonids (*Oncorhynchus* spp.) (U.S. Army Corps of Engineers 1992a). However, possible adverse effects of a drawdown include changes in the benthic invertebrate community, which has been found to be an important constituent of juvenile salmonid diet in the Columbia River system (Becker 1973, Kim et al. 1986, Muir and Emmett 1988, Muir and Coley 1996).

Several drawdown scenarios have been proposed for selected Columbia and Snake River reservoirs. Each scenario involves the operation of reservoirs at or below minimum operating pool (MOP) for several months each year. Any extended drawdown at or below MOP would be an unprecedented event in the operation of these reservoirs and would have undetermined impacts on reservoir and associated riverine ecosystems.

For Lower Granite Reservoir, a drawdown to depths of 10, 13, and 16 m below MOP during the spring and summer juvenile salmonid outmigration periods (April-August) has been suggested. A drawdown to depths of 33.5 to 35 m below MOP to simulate a free-flowing river has also been suggested. Such a drawdown would substantially reduce existing shallow-water habitat (U.S. Army Corps of Engineers 1992a). Furthermore, the biological productivity of current deepwater habitats may not sufficiently replace the existing

shallow-water habitats after a drawdown, and this may result in a decrease in available food for juvenile salmonids.

To better understand possible changes that may occur as a result of a drawdown, limnological studies commenced in 1994 to document pre-drawdown conditions (Ledgerwood et al. 1996, Juul et al. 1997, Bennett et al. 1997). One objective of these studies covered by this report was to document the species composition and densities of benthic invertebrates inhabiting soft-substrate, shallow-water habitats.

Benthic invertebrate samples were collected from three soft-substrate, shallow-water sampling areas within the reservoir. Sampling areas were located near the upstream end of the reservoir at River Kilometer (RKm) 212 near Silcott Island, near mid-reservoir at RKm 193 near Centennial Island, and near the terminus of the reservoir at RKm 177 about 3 km upstream from Offfield Landing. At each sampling area, four benthic invertebrate samples were collected from three depth contours (3, 9, and 18 m). Between March 1994 and October 1995, benthic invertebrate samples were collected monthly, except in April and December 1994 and February 1995, using a Ponar grab to sample about 0.05 m². A total of 647 benthic invertebrate samples was collected, and more than 4,000 hours were required to process all the samples.

A total of 76 taxa/categories was identified from the processed samples. Oligochaeta, Chironomidae larva (order Diptera), and to a lesser extent Bivalvia were numerically dominant in both density and frequency of occurrence at all sampling areas and depths; overall these 3 taxa/categories, which were found throughout the reservoir, comprised 93% of all organisms enumerated. Mean densities of all benthic invertebrates, except oligochaetes,

tended to decrease with depth in all sampling areas. Oligochaete densities did not decrease with depth except near Silcott Island.

There was no strong seasonal trend in oligochaete or chironomid larva density at any sampling area; overall mean densities were 6,231 and 845/m² (data pooled by date, sampling area, and depth), respectively. Bivalves (generally immature) were the third most abundant benthic invertebrates (overall mean = 235/m²) and at Offield accounted for about 75% of all bivalves recovered. Amphipods were the predominant benthic crustaceans in the reservoir (overall mean = 14/m²). Amphipods were present sporadically at all areas and depths sampled and five species were identified.

At the upper reservoir sampling area near Silcott Island, densities of oligochaetes and chironomid larvae decreased with depth, a trend possibly correlated with the percentages of sandy and fine sediment composition in the area. In contrast, at the mid-reservoir sampling area at Centennial Island, densities of oligochaetes and chironomid larvae were not apparently correlated with depth, and the percentages of fines at different depths were fairly consistent. The lower reservoir sampling area at Offield was the only area with gravel substrate, a component which may have been related to increased bivalve densities in that area.

CONTENTS

EXECUTIVE SUMMARY	iii
INTRODUCTION	1
METHODS	2
Study Area	2
Collection	3
Sample Processing	6
Sediment Analysis	7
Data Analysis	8
Potential Drawdown Effects	8
RESULTS	9
Benthic Invertebrate Community	9
Dominant Benthic Invertebrates	13
Oligochaetes	13
Chironomid larvae	17
Bivalves	17
Other Benthic Invertebrates	20
Aquatic Insects	20
Crustaceans	23
Sediment Samples	24
DISCUSSION	25
CONCLUSIONS	29
RECOMMENDATIONS	30
REFERENCES	33
APPENDICES	39
Appendix A	41
Appendix B	45
Appendix C	57
Appendix D	85

INTRODUCTION

Historical estimates of adult salmon abundance in the Columbia River basin before hydroelectric dams were built range from 8 to 16 million, but since then estimates range from 2 to 2.5 million (U.S. Army Corps of Engineers 1992a). Snake River sockeye salmon (*Oncorhynchus nerka*) was listed as endangered in 1991 and spring/summer and fall chinook salmon (*O. tshawytscha*) were listed as threatened in 1992 under the Federal Endangered Species Act of 1973. A "Salmon Summit" was held in 1990 and 1991 between regional fishery and power agencies, tribes, and river users to develop a restoration plan for declining salmon runs. At the summit, one suggestion was to draw the river water level down during juvenile salmon migration, which would theoretically increase downstream water velocity and thus decrease travel time for salmon smolts migrating to the ocean. A 1-month test drawdown of Lower Granite Reservoir occurred during March 1992.

In 1994, personnel of the Coastal Zone and Estuarine Studies (CZES) Division of the National Marine Fisheries Service and other entities began limnological investigations to assess possible impacts of a drawdown on Lower Granite Reservoir (Ledgerwood et al. 1996, Juul et al. 1997, Bennett et al. 1997). This report covers CZES limnological investigations dealing with soft-substrate, shallow-water benthic invertebrate populations sampled between March 1994 and October 1995.

Benthic invertebrates are an important food source for juvenile salmon in the Columbia River system (Becker 1973, Kim et al. 1986, Muir and Emmett 1988, Muir and Coley 1996). The overall importance of soft-substrate, shallow-water habitat to outmigrating juvenile salmonid population is unclear. Bennett et al. (In prep.) reported that age-0 fall

chinook salmon had higher abundance over sand and mud/sand substrates in Lower Granite Reservoir. If drawdown were to reduce the availability of such habitat, then it could be detrimental to recovery efforts for these endangered fish. The goal of this study was to document the species composition, spatial variation, and temporal distribution of benthic invertebrates inhabiting soft-substrate, shallow-water habitats in the reservoir during the pre-drawdown conditions and to compare these with results during or following a possible drawdown.

METHODS

Study Area

Lower Granite Reservoir was created in 1975 when Lower Granite Dam was constructed for hydroelectric power, navigation, and irrigation (U.S. Army Corps of Engineers 1992b). The reservoir is located in southeastern Washington and western Idaho near Clarkston, Washington, and Lewiston, Idaho. The reservoir extends 61.8 km from the dam to Asotin, Washington on the Snake River and 7.3 km upstream from the confluence of the Snake and Clearwater Rivers on the Clearwater River (U.S. Army Corps of Engineers 1992b). Following construction of the dam, the hydrography of this segment of the Snake River changed from free-flowing river to a pool with a maximum depth of 35 m (Dorband 1980).

Collection

Three soft-substrate, shallow-water areas in Lower Granite Reservoir were chosen for sampling (Fig. 1). These areas were at River Kilometer (RKm) 212 near Silcott Island, at RKm 193 near Centennial Island, and at RKm 177 near Offield Landing. The sampling area near Silcott Island was located about 39 km upstream from Lower Granite Dam and about 11 km downstream from the confluence of the Snake and Clearwater Rivers at Lewiston, Idaho and Clarkston, Washington. Centennial Island was created near the middle of the reservoir in 1989 as a result of dredging activity by the U.S. Army Corps of Engineers (U.S. Army Corps of Engineers 1992b) and is about 20 km upstream from the dam. The Offield sampling area is about 4 km upstream from Lower Granite Dam (RKm 173).

From March 1994 through October 1995, benthic invertebrate samples were collected monthly, except in April and December 1994 and February 1995. A Ponar grab (Word 1976) was used to collect samples of about 0.05 m² along four transects perpendicular to shore at 3-, 9-, and 18-m depths (Fig. 2). A total of 12 samples was taken from each sampling area each month. Although the 18-m sample depth lies beyond the shallow-water zone defined for this study, samples from this depth were necessary to determine conditions in the existing deepwater zone, which will become a shallow-water zone after a drawdown of up to 16 m. At Centennial Island, two samples from the 3-m depth were collected from the inside passage along the island: one from the middle upstream transect and the other from the middle downstream transect (Fig. 2).

Samples were initially washed in the field through a 0.5-mm sieve (U.S.A. Standard Testing Sieve no. 35) and then preserved in a solution of about 10% buffered formalin. The




Figure 1. Location of Lower Granite Reservoir and locations of three selected soft-substrate, shallow-water sampling areas.




Figure 2. Schematic drawings (not to scale) of the benthic invertebrate sampling areas in Lower Granite Reservoir in 1994-1995. Four transects perpendicular to shoreline were established in each sampling area and numbered boxes indicate a station and depth (m). Boundaries of sampling areas \pm SE are shown based on hand-held Global Positioning System recordings.

preservative contained rose bengal which stained organisms to speed processing in the laboratory.

Sample Processing

In the laboratory, each sample was washed again with tap water through a 0.5-mm sieve (U.S.A. Standard Testing Sieve no. 35) to remove formalin and fine sediments prior to processing. Dissecting microscopes with up to 40X magnification were used to sort through the sieved residue, and we recovered at least 90% of all organisms, except eggs and cladocerans. Organisms removed from each sample were stored in a small vial with ethanol and a drop of glycerin. Eggs were not included in the recovery process because it was not feasible to determine whether they were of invertebrate or vertebrate origin. Cladocerans are mainly planktonic, not benthic invertebrates; hence, they were not included.

Generally, identifications were made to the lowest practical taxon (Barnard 1969, Smith and Carlton 1975, Borror et al. 1976, Pennak 1978, Bousfield 1979, Rudy and Rudy 1983, Barnes 1987). Insects were further identified by life history stage (larva/nymph, pupa, and adult). Only heads of fragmented organisms were counted. Fragmented oligochaetes were commonly found and often had phenotypically similar ends, rather than a distinct head. Therefore, in the case of fragmented oligochaetes, parts of similar widths and with a terminal end were paired together and counted as one individual.

For each benthic invertebrate sample, we recorded the identification and enumeration of the organisms recovered, the sample's labelling information, the processor's name, the date(s) the sample was processed, and the number of processing hours (Appendix Fig. D1).

To ensure the removal of at least 90% of all organisms except eggs and cladocerans from processed samples, a minimum of 10% of all samples were verified.

Three to five samples collected each sampling period were large in volume (1-2 L after sieving). These large samples required an estimated 20 to 40 hours each to process using standard procedures. To use laboratory time more efficiently, a 4-L capacity plankton splitter was utilized to split a large sample into 6 equivalent subsamples (Appendix A). Two of the six subsamples were subsequently processed and the counts used to estimate total tallies for the entire sample; 56 large samples were subsampled and processed using this procedure.

Sediment Analysis

In July 1995, a sediment sample was collected from each of the 36 benthic invertebrate sampling stations. Sediment analyses for particle grain size, soil classification, and percent volatile solids were conducted under contract to the U.S. Army Corps of Engineers North Pacific Division Materials Laboratory, Troutdale, Oregon. We used these analyses to evaluate possible differences in sediment composition as related to benthic invertebrate populations among transects at each sampling area.

Data Analysis

The National Oceanographic Data Center (NODC) taxonomic codes (version 7.0) were assigned to each organism based on its taxonomic classification.¹ Taxonomic information is contained in the hierarchy of each code and allows data for organisms to be grouped and summed into a desired taxonomic classification. Computer programs were developed to allow organisms to be grouped into five possible taxonomic levels for data analysis; for example, *Hexagenia* spp., Caenidae nymphs, and other Ephemeroptera nymphs could be combined into the Ephemeroptera order, and total tallies for this insect order could be compared with those of other insect orders. Mean densities were calculated for each grouped or ungrouped taxon/category for comparisons among sampling areas, depths, and dates. Density was expressed as number of organisms/m². Although terrestrial, planktonic, epibenthic, and benthic organisms were found in our benthic invertebrate samples, analysis focused on benthic and epibenthic invertebrates (Appendix Tables B1 and B2). We further focused our presentations in this report on the three most abundant taxa/categories found in the sampling areas. In addition, benthic invertebrates grouped into subphyla Insecta and Crustacea were further analyzed at the subphylum taxonomic level.

Potential Drawdown Effects

Drawdown was removed as an option for the Lower Granite Reservoir in the fall of 1995. As a result, sampling of the pre-drawdown condition was discontinued. Lacking

¹ National Oceanographic Data Center, NOAA/NESDIS E/OC1, SSMC3, Room 4649, 1315 East-West Highway, Silver Spring, MD 20910-3282

information on a drawdown or post-drawdown condition, we could not conduct detailed statistical analyses of the benthic invertebrate data for this report.² Temporal changes in the pre-drawdown soft-substrate, shallow-water benthic invertebrate community of Lower Granite Reservoir between March 1994 and October 1995 are presented.

RESULTS

Benthic Invertebrate Community

A total of 647 benthic invertebrate samples was collected from our 3 sampling areas during 1994-95. These required an average of 6.5 hours per sample for processing (over 4,000 hours total). All samples were processed within 2 years of collection. Over 250,000 organisms were recovered and enumerated during sample processing (Appendix C). Details of taxonomic groupings and electronic formats of the data are available upon request.

Overall, a total of 76 taxa/categories was found within the sampling areas. Of the sampling areas, Silcott Island had the highest density of organisms (12,578/m²) compared to Centennial Island (7,069/m²) and Offield (3,842/m²). Each taxon/category of organisms found in the reservoir with its associated NODC code is listed in Table 1. Most of the benthic invertebrates recovered were in the Oligochaeta, Insecta, Bivalvia, and Crustacea taxa/categories (Fig. 3).

² The data presented for the pre-drawdown condition would benefit from statistical analyses (cluster analyses, diversity indices, and possibly other tests) but cancellation of the study compromised our ability to complete these analyses at this time.

Table 1. Taxa/categories found in benthic invertebrate samples collected from Lower Granite Reservoir, 1994-1995.

Taxon/category	NODC code ^a	Taxon/category	NODC code ^a
Platyhelminthes		Insecta	620000000000
Turbellaria	390100000000	Insecta adult	620000000093
Nemertea	430000000000	Coleoptera	630200000000
Nematoda	470000000000	Coleoptera adult	630200000093
Nematomorpha	480000000000	Coleoptera larvae	630200000091
Mollusca		Elmidae adult	631604000093
Gastropoda	510000000000	Collembola	620800000000
Archaeogastropoda	510200000000	Ephemeroptera nymph	621500000091
Bivalvia	550000000000	Caenidae nymph	621802000091
Annelida		<i>Caenis</i> spp.	621802020000
Oligochaeta	500300000000	<i>Ephemera</i> spp.	622003020000
Polychaeta	500100000000	<i>Hexagenia</i> spp.	622003030000
Hirudinea	501200000000	Leptophlebiidae nymph	621701000000
Crustacea		Hemiptera	627100000000
Cladocera	610800000000	Hemiptera adult	627100000093
Leptodoridae	610906000000	Homoptera adult	628200000093
<i>Leptodora kindtii</i>	610906010100	Formicidae adult	657307000093
Ostracoda	611000000000	Lepidoptera larvae	642000000091
Gammaridae	616921000000	<i>Sialis</i> spp.	640601010000
<i>Corophium</i> spp.	616915020000	Plecoptera	625100000000
<i>Corophium salmonis</i>	616915020900	Plecoptera adult	625100000093
<i>Corophium spinicorne</i>	616915021500	Plecoptera nymph	625100000091
<i>Ramellogammarus</i>	616921460200	Psocoptera	625600000000
<i>oregonensis</i>		Thysanoptera adult	626800000093
<i>Ramellogammarus</i>	616921460100	Diptera	648100000000
<i>ramellus</i>		Diptera adult	648100000093
<i>Hyaella azteca</i>	616923040100	Diptera larvae	648100000091
Isopoda	615800000000	Diptera pupae	648100000092
<i>Porcellio</i> spp.	616604040000	Chironomidae adult	648933000093
Mysidacea	615100000000	Chironomidae larvae	648933000091
Copepoda	611700000000	Chironomidae pupae	648933000092
Cyclopoida	612000000000	Chironominae pupae	648959000092
Harpacticoida	611900000000	Orthoclaadiinae pupae	648956000092
Calanoida	611800000000	Tanypodinae pupae	648938000092
Chelicerata		Ceratopogonidae larvae	648920000091
Araneae	591100000000	Culicidae adult	648906000093
Prostigmata	592900000000	Simuliidae larva	648915000091
Ixodides	592800000000	Tanyderidae larvae	648804000091

Table 1. Continued.

Taxon/category	NODC code	Taxon/category	NODC code*
Insecta continued		Insecta continued	
Orthoptera adult	623100000093	Isoptera	624600000000
Trichoptera adult	641800000093	Miscellaneous	
Trichoptera larvae	641800000091	Eggs (unidentified)	
Psychomyiidae larvae	641803000091	Unidentified	

- * The NODC codes listed are taxonomic numerical codes assigned by the National Oceanographic Data Center. For insects, the last two digits were used to indicate the life history stage. Except at the genus level, 91, 92, and 93 were used to represent the larval/nymph, pupa, and adult stages respectively.




Figure 3. Relative composition of major benthic taxa found in three soft-substrate, shallow-water sampling areas (pooled data) of Lower Granite Reservoir, 1994-95.

Dominant Benthic Invertebrates

Three taxa/categories were found in the reservoir: Oligochaeta, Chironomidae larvae, and Bivalvia, and these comprised 93% of all organisms enumerated (Table 2). Oligochaetes and chironomid larvae were also numerically dominant in each sampling area. Bivalves were the third most abundant benthic invertebrates at Centennial Island and Offfield, but not at Silcott Island.

Oligochaetes--Oligochaeta was the most abundant taxon/category at all sampling areas and appeared to increase in abundance from 1994 to 1995 at all depths at each sampling area (Fig. 4). Oligochaetes were most abundant at Silcott Island, especially at the 3-m depth. At Silcott Island, respective mean oligochaete densities at the 3-, 9-, and 18-m depths were as follows: 18,220, 10,865 and 4,819/m². This apparent decrease in oligochaete density with depth at Silcott Island was consistent throughout the study period. At all sampling areas, temporal changes in oligochaete density were not apparent.

Unlike at Silcott Island, densities of oligochaetes at Centennial Island and Offfield were higher along the 18-m depth contour (Fig. 4). Respective mean oligochaete densities at the 3-, 9-, and 18-m depths were as follows: 5,891, 3,166, and 6,565/m² at Centennial Island and 1,803, 1,698, and 3,047/m² at Offfield.

There were apparent differences in oligochaete densities among transects at each depth in each sampling area. These differences appeared greatest at the 3-m depth at Silcott Island and the 9- and 18-m depths at Centennial Island (Appendix Fig. D2-D3). Of the three sampling areas, Offfield had the least differences in oligochaete densities among transects at

Table 2. Mean densities (no./m²) and standard deviations (SD) of benthic taxa/categories by year and sampling area in the Lower Granite Reservoir.

Taxon/category	Silcott Island RKm 212				Centennial Island RKm 193				Offfield RKm 177			
	1994		1995		1994		1995		1994		1995	
	No./m ²	SD	No./m ²	SD	No./m ²	SD	No./m ²	SD	No./m ²	SD	No./m ²	SD
Platyhelminthes												
Turbellaria	3	13	59	401	4	24	3	13	1	8	2	8
Nemertea	15	106	10	51	14	61	15	35	9	39	4	24
Mollusca												
Gastropoda	1	6	1	6	<1	2	0	0	<1	2	0	0
Bivalvia	10	22	36	75	123	304	111	297	461	796	399	660
Annelida												
Oligochaeta	8,523	9,442	14,096	15,394	4,134	3,894	6,281	5,593	1,462	1,915	2,904	3,147
Polychaeta	5	22	4	19	<1	2	<1	3	2	6	5	18
Hirudinea	1	3	1	6	1	4	<1	4	<1	3	<1	3
Crustacea												
Ostracoda	32	64	24	53	22	63	26	50	49	130	24	46
Unidentified Gammaridae ^a	0	0	0	0	0	0	0	0	<1	4	<1	4
<i>Corophium</i> spp.	4	29	2	13	1	5	2	8	20	101	55	283
<i>Ramellogammarus</i> spp.	1	3	0	0	1	6	0	0	<1	3	0	0
<i>Hyalella azteca</i>	0	0	0	0	0	0	0	0	<1	2	0	0
Isopoda	<1	2	0	0	0	0	0	0	1	7	0	0
Mysidacea	0	0	0	0	1	10	0	0	0	0	0	0
Copepoda ^b	42	113	32	81	68	201	54	151	39	116	32	109

Table 2. Continued.

Taxon/category	Silcott Island RKm 212				Centennial Island RKm 193				Offield RKm 177			
	1994		1995		1994		1995		1994		1995	
	No./m ²	SD	No./m ²	SD	No./m ²	SD	No./m ²	SD	No./m ²	SD	No./m ²	SD
Chelicerata												
Prostigmata	4	11	15	72	24	67	19	41	45	75	40	66
Insecta												
Coleoptera ^a	1	4	1	6	0	0	<1	2	<1	3	1	4
Ephemeroptera nymph	2	8	5	16	5	15	9	32	80	243	39	83
Lepidoptera larvae	0	0	2	14	0	0	0	0	0	0	0	0
Megaloptera larvae	0	0	0	0	0	0	0	0	<1	2	0	0
Plecoptera nymph	1	3	4	26	1	4	<1	2	0	0	0	0
Unidentified Diptera larvae	<1	2	0	0	<1	2	0	0	0	0	0	0
Unidentified Diptera pupae	1	6	4	29	0	0	0	0	<1	4	0	0
Chironomidae larvae	917	1,132	727	924	1,014	925	846	755	916	1,116	649	729
Chironomidae pupae	7	18	4	16	13	31	5	17	16	47	5	12
Ceratopogonidae larvae	<1	2	<1	2	<1	2	<1	2	2	9	<1	2
Tanyderidae larvae	0	0	1	6	0	0	0	0	0	0	0	0
Trichoptera larvae	2	7	5	17	5	18	2	6	3	10	3	12

^a Unidentified Gammaridae exclusive of *Corophium* spp.^b Includes harpacticoids and unidentified copepods.^c Includes all life history stages.




Figure 4. Densities (no./m²) of Oligochaeta by depth at three sampling areas in Lower Granite Reservoir, 1994-1995.

all depths with the greatest difference observed in 1995 along the 18-m depth contour (Appendix Fig. D4).

Chironomid larvae--Chironomid larvae ranked second in abundance at each sampling area. At Silcott Island, mean chironomid larva densities decreased with depth (Fig. 5). Respective mean chironomid larva densities at the 3-, 9-, and 18-m depths were as follows: 1,191, 905, and 394/m² at Silcott Island; 943, 953, and 894/m² at Centennial Island; and 986, 544, and 816/m² at Offfield.

Differences among transects for each sampling depth in each sampling area were greater for chironomid larvae than for oligochaetes. At Silcott Island, differences in chironomid larva densities among transects were small at the 18-m depth and large at the 3- and 9-m depths (Appendix Fig. D5). At Centennial Island, differences in chironomid larva densities among transects were similar at all sample depths (Appendix Fig. D6). At Offfield, the greatest variation of chironomid larva densities among transects was along the 3-m depth contour (Appendix Fig. D7).

Bivalves--Bivalves were the third most abundant organisms in the sampling areas (data pooled by sampling areas, date, and depths). Offfield had the highest abundance of bivalves, accounting for about 75% of bivalves in all three sampling areas (Fig. 6). Respective mean bivalve densities at the 3-, 9-, and 18-m depths were as follows: 23, 18, and 28/m² at Silcott Island; 321, 26, and 4/m² at Centennial Island; and 665, 378, and 248/m² at Offfield. At Offfield, mean bivalve densities were higher from October 1994 to May 1995 (mean = 665/m²) at all sampling depths than during other months, and mean densities were lowest from June to September (mean = 294/m²).




Figure 5. Densities (no./m²) of Chironomidae larvae by depth at three sampling areas in Lower Granite Reservoir, 1994-1995.




Figure 6. Densities (no./m²) of *Bivalvia* by depth at three sampling areas in Lower Granite Reservoir, 1994-1995.

Differences in bivalve densities among transects for each sampling depth were not apparent except at the 3-m depth contour at Offfield (Appendix Figs. D8-D10). At Silcott Island and Centennial Island, bivalve densities were generally less than $0.5/\text{m}^2$ in our samples at all depth contours. At the 9- and 18-m depth contours at Offfield, though bivalve densities often exceeded $0.5/\text{m}^2$, there was no obvious pattern of difference among transects.

Other Benthic Invertebrates

Aquatic Insects--Aquatic insects comprised 11% of all organisms enumerated. Included in decreasing order of abundance were the following seven orders of aquatic insects: Diptera, Ephemeroptera, Trichoptera, Plecoptera, Coleoptera, Lepidoptera, and Megaloptera (Table 3). Dipteran and ephemeropteran insects were generally abundant and found at all sampling areas and depths throughout the study period. The remaining insect orders had densities generally less than $10/\text{m}^2$ and were present only sporadically during the study period (Table 3).

The most abundant insects were dipterans (Table 3), which included the Chironomidae larvae and pupae, Ceratopogonidae larvae, and Tanyderidae larvae taxa/categories. Mean dipteran densities ranged from 20 to $3,230/\text{m}^2$ for all sampling areas and depths, with mean densities pooled by depth at Silcott Island, Centennial Island, and Offfield of 832, 940, and $794/\text{m}^2$, respectively (Fig. 7). There was large seasonal variation in dipteran densities at all sampling areas and depths.

Ephemeroptera (mayfly nymphs) was the second most abundant order of aquatic insects (Table 3). Temporally, mean mayfly nymph densities ranged from 0 to $725/\text{m}^2$, with highest densities at Offfield (Appendix Table C). Mean mayfly nymph densities at Silcott

Table 3. Mean density (no./m²) of Insecta orders collected at three soft-substrate, shallow-water sampling areas in Lower Granite Reservoir, 1994-1995. Data from all sampling dates are pooled for each sampling area and water depth.

Sampling area	Depth (m)	Coleoptera	Diptera	Ephemeroptera	Lepidoptera	Megaloptera	Plecoptera	Trichoptera
Silcott Island	3	1	1,184	7	3	0	2	6
	9	1	914	3	0	0	4	3
	18	0	398	1	0	0	1	1
Centennial Island	3	0	961	16	0	0	0	9
	9	0	959	2	0	0	<1	<1
	18	<1	899	3	0	0	1	1
Offfield	3	<1	1,013	141	0	0	0	6
	9	<1	549	35	0	<1	0	2
	18	1	820	3	0	0	0	<1




Figure 7. Densities (no./m²) of Diptera by depth at three sampling areas in Lower Granite Reservoir, 1994-1995.

Island, Centennial Island, and Offfield were 4, 7, and 59/m², respectively. At all sampling areas, the highest mayfly nymph density was at the 3-m depth, and mayfly nymphs were rare at the 18-m depth.

Crustaceans--Five taxa of crustaceans were found that can be included as benthic invertebrates. These taxa were as follows, in decreasing order of abundance: Copepoda, Ostracoda, Amphipoda, Isopoda, and Mysidacea (Appendix Table B1). Copepods and ostracods were the two most abundant crustaceans and were found at each sampling area and depth (Appendix Table B3). Isopods and mysids were rare (Appendix Table B3). Isopods were present during three sampling periods and mysids were present once.

Mean copepod densities pooled by depths ranged from 0 to 390/m² (Appendix Table B3). Respective mean copepod densities at the 3-, 9-, and 18-m depths were as follows: 71, 25, and 14/m² at Silcott Island; 38, 92, and 54/m² at Centennial Island; and 24, 28, and 54/m² at Offfield. At Silcott Island 9- and 18-m depths and at the Centennial Island 18-m depth, copepod densities peaked in August 1994. At Offfield 9- and 18-m depths and at the Centennial Island 3-m depth, copepod densities peaked in November 1994. During other months, mean copepod densities were generally less than 100/m².

Ostracods were next in abundance to copepods, and in samples from some months, ostracods had higher densities than copepods (Appendix Table B3). Respective mean ostracod densities at Silcott Island, Centennial Island, and Offfield were 28, 24, and 37/m². Generally, mean ostracod densities were less than 100/m² throughout the study.

Although amphipods were sporadically present at all areas and depths sampled, they were higher in density at times than the copepods and ostracods (Appendix Table B3). For

example, at Offfield in April 1995, amphipod densities at the 3- and 9-m depths were 620 and 430/m², respectively. At least five species of amphipods were identified: *Corophium salmonis*, *C. spinicorne*, *Ramellogammarus oregonensis*, *R. ramellus*, and *Hyaella azteca*. The predominant genus of amphipods present in the study areas was *Corophium*. Overall, the mean amphipod density was 14/m².

Sediment Samples

There were apparent differences in sediment composition among sampling areas, between depths within sampling areas, and along transects within sampling areas (Appendix Table B4-B5). Percentages of gravel (grain sizes 75 to 4.75 mm) were zero at all stations at Silcott Island and Centennial Island, whereas at Offfield, percentages of gravel in the sediments ranged from 0 to 38.8. Percentages of sand (grain sizes 4.75 to 0.074 mm) ranged from 12.9 to 94.0 at Silcott Island, from 4.3 to 79.6 at Centennial Island, and from 6.2 to 58.0 at Offfield. At Silcott Island, the percentages of sand were highest at the 18-m depth, while at Centennial Island they were highest at the 3-m depth. Percentages of sand at Offfield appeared more variable than at the other two sampling areas.

Percentages of fines (grain sizes < 0.074 mm) ranged from 6.0 to 87.1 at Silcott Island, from 20.4 to 95.7 at Centennial Island, and from 54.6 to 93.8 at Offfield. At Silcott Island, percentages of fines were higher at the 3-m depth than at the 9- and 18-m depths. At Centennial Island and Offfield, percentages of fines were higher at the 9- and 18-m depths than at the 3-m depth. The median grain size ranged from 0.03 to 0.18 mm at Silcott Island, from 0.02 to 0.28 mm at Centennial Island, and from 0.01 to 0.36 mm at Offfield.

Percentages of silt/clay ranged from 5.4 to 84.5 at Silcott Island, from 20.3 to 95.0 at Centennial Island, and from 32.4 to 90.6 at Offield. Centennial Island had the highest percentages of silt/clay, especially at the 9- and 18-m depths. Percentages of volatile solids ranged from 0.4 to 48.3 at Silcott Island, from 2.1 to 15.8 at Centennial Island, and from 3.8 to 10.0 at Offield. There was no consistent pattern of sediment composition among transects within sampling areas.

DISCUSSION

Benthic invertebrate fauna in the three selected soft-substrate, shallow-water habitats of Lower Granite Reservoir during 1994-95 was numerically dominated by oligochaetes and to a lesser extent, chironomid larvae and bivalves; a distribution similar to results reported for 1976-77 (Dorband 1980). Mean oligochaete densities in our sampling areas ranged from 1,698 to 18,220/m² and the mean chironomid larva density was about 850/m² with a peak density over 3,000/m². Dorband (1980) reported that densities for both oligochaetes and chironomids were similar in 1976 and 1977 (500 to 13,000/m² and 492 to 1,292/m², respectively). Insect orders recovered from our benthic invertebrate samples included all of those reported by Dorband (1980). Bennett et al. (1990, 1993) found that oligochaetes and chironomids dominated the benthic fauna during June, October, and December 1988, July 1989, and September 1991. They reported ratios of 45% oligochaetes to 55% chironomids in standing crop estimates for the reservoir and a mean numerical density for chironomids of about 500/m² with chironomid densities occasionally reaching at 2,000/m². It appears that the

benthic invertebrate community in Lower Granite Reservoir has been fairly stable for the past 20 years.

We chose to index our benthic invertebrate data analyses by a single variable, numerical density. Multiple variables like number, weight, and frequency of occurrence in multivariant analysis are difficult to interpret, require extra effort to collect, and increase sensitivity to violations of assumptions (Macdonald and Green 1983). Dorband (1980) reported numerical densities and frequency of occurrence for organisms recovered from benthic invertebrate samples collected from Lower Granite, Little Goose, and Ice Harbor Reservoirs. Bennett et al. (1993) reported both density and weights of chironomids in Lower Granite Reservoir but did not report numerical data for oligochaetes due to problems associated with counting fragmented oligochaetes. We also had difficulty with enumerating fragmented oligochaetes; therefore, our reported oligochaete densities are possibly biased high. However, we believe the degree of possible bias would not compromise comparisons required for assessment of a possible drawdown.

We did not attempt further taxonomic identification of oligochaetes because it was considered beyond the scope of this study. For accurate identification of oligochaetes to genera and species, it is necessary to serially section and mount fresh specimens on slides for examination of the anatomical arrangement (Pennak 1978). However, Dorband (1980) reported that most oligochaetes in Lower Granite Reservoir were identified as either *Tubifex tubifex* or *Limnodrilus hoffmeisteri*. Changes in genera or species of oligochaetes or other benthic invertebrates could be important to fully understand possible impacts of a drawdown on the benthic invertebrate community in the reservoir.

The abundance of some taxa/categories in the upper reservoir may be due in part to the sharp, wide turn of the reservoir at Silcott Island, where the westerly and downstream water flow turns sharply northwest and continues through the steep-sloped canyon to the dam. Silcott Island probably receives more turbid and silty waters from the convergence of the Clearwater and Snake Rivers because this area is closest to the riverine-to-reservoir transition than the two downstream sampling areas. Hence, the higher percentages of sand and fines at Silcott Island provides improved habitat for oligochaetes and chironomids.

Differences in species composition and abundance and temporal changes in densities of benthic invertebrates at our sampling areas were undoubtedly affected by differences in sediment composition. The predominantly sand and fine sediment composition in the soft-substrate, shallow-water habitats we studied probably contributed to the dominance of oligochaetes and chironomid larvae. Some freshwater species of oligochaetes prefer mud and silt for burrow-making and feed on organic matter and bacteria present in those sediments (Brinkhurst and Cook 1974, Barnes 1987, Brinkhurst and Gelder 1991).

Chironomid larvae have been found in almost all aquatic habitats, including mud, sand, and rocks (Roback 1974, Pennak 1978). Aquatic insects, in addition to chironomid larvae, have habitat preferences which also depend on sediment types. For example, some mayfly nymphs are more adapted to gravel than other invertebrates (Roback 1974, Pennak 1978). This adaptation to gravel may explain the higher mayfly nymph densities at Offfield, the only sampling area with gravel. However, other mayfly nymphs we found, such as *Hexagenia* spp. and *Caenis* spp., prefer soft mud bottoms and slow, silty water areas (Roback 1974, Pennak 1978). Other factors possibly affecting aquatic insect densities include

emergence of subadults and predation. Populations of bivalves are reportedly higher in habitats where the benthos is stabilized with gravel and sand, rather than shifting sand and mud (Pennak 1978). The gravel and sand substrate at Offield may contribute to higher bivalve densities in that sampling area. Amphipods, the predominant benthic crustaceans in our sampling areas, generally live in and among debris and stones and burrow into sediments (Pennak 1978, Barnes 1987). Amphipods were present at low densities in all sampling areas (mean = 14/m²), but the reported densities may also be affected by concentration of carbonate in the water, dissolved oxygen concentration, or other biotic and abiotic factors (Pennak 1978).

The proposed drawdown would expose much existing shallow-water habitat of the reservoir, as was discovered during a drawdown test in March 1992 (U.S. Army Corps of Engineers 1992a). This exposure may have important impacts on the diets of juvenile salmonids and other fishes if there was a net loss of such habitat following a drawdown.

Insects, primarily emerging subadult chironomids, are an important prey of juvenile salmonids in the Columbia River system (Becker 1973, Kirn et al. 1986, Muir and Emmett 1988, Muir and Coley 1996). Chironomid larva also constitutes one of major food resources for juvenile and adult fishes in freshwater habitats (Pennak 1978). In their analysis of stomach contents of migrating juvenile chinook salmon passing through Lower Granite Dam, Muir and Coley (1996) reported six orders of aquatic and terrestrial insects, with chironomids as the dominant prey. Each of the insect orders observed by Muir and Coley (1996) was also recovered from our benthic invertebrate samples. Therefore, it is possible that temporal densities of insects in Lower Granite Reservoir were affected by predation as well as

emergence of subadults, and that loss of benthic fauna could affect food resources of juvenile salmonids.

Oligochaetes, though the most abundant benthic invertebrates in the reservoir, have not to our knowledge been reported as a major diet constituent for juvenile salmonids. Muir and Emmett (1988) reported that bivalves (*Corbicula manilensis*) were rarely found in stomachs of juvenile salmonids. Hence, neither oligochaetes nor bivalves, which were among the most abundant benthic invertebrates in our sampling areas, are likely important dietary components for salmonids. Crustaceans, except amphipods, were reportedly less important than insects as prey of juvenile salmonids in the Columbia River system (Becker 1973, Kirn et al. 1986, Muir and Emmett 1988, Muir and Coley 1996). Amphipods, especially *Corophium* spp., were reported to be the dominant prey of juvenile salmonids at Bonneville Dam (Muir and Emmett 1988). At Lower Granite Dam, outmigrating juvenile chinook salmon prey included amphipods (*Corophium spinicorne*) (Muir and Coley 1996). The potential contribution of these benthic invertebrates to juvenile salmonids in Lower Granite Reservoir should not be overlooked.

CONCLUSIONS

- 1) The selected soft-substrate, shallow-water areas sampled during 1994-95 were numerically dominated by oligochaetes, aquatic insects, primarily chironomid larvae, and bivalves. Although 76 taxa/categories of benthic invertebrates were present in these areas, mean densities at each sampling area were <13,000 organisms/m².

- 2) Densities of benthic invertebrates decreased from the upper reservoir sampling area near Silcott Island (mean = 12,578/m²) to the mid-reservoir sampling area at Centennial Island (mean = 7,069/m²) and decreased further from Centennial Island to the lower reservoir sampling area near Offield Landing (mean = 3,842/m²). These density differences were possibly related to differences in sediment composition among sampling areas.
- 2) At present reservoir pool levels, soft-substrate, shallow-water habitats are rare due to the steep-sloped canyon of the reservoir. The overall impact of a drawdown on benthic invertebrates present will depend in part on the net gain or loss of these habitats following a drawdown.

RECOMMENDATIONS

- 1) Activities affecting production of food resources for juvenile salmonids could affect restoration efforts for threatened or endangered Snake River salmon. Benthic invertebrates are an important dietary component of juvenile salmonids and activities, such as drawdown, which could affect food availability should be carefully monitored.
- 2) Statistical analyses to examine possible differences in composition and abundance of benthic invertebrates among dates, depths, transects, and sampling areas are needed. Analyses to evaluate species diversity and relationships between benthic invertebrates

and other limnological parameters monitored during this pre-drawdown period are also recommended.

REFERENCES

- Barnard , J. L. 1969. The families and genera of marine gammaridean Amphipoda. Smithsonian Institution Press, City of Washington. U. S. Nat. Mus. Bull. 271, 535 p.
- Barnes, R. D. 1987. Invertebrate zoology, fifth edition. Saunders College Publishing. U.S.A., 893 p.
- Becker, C. D. 1973. Food and growth parameters of juvenile chinook salmon, *Oncorhynchus tshawytscha*, in central Columbia River. Fish. Bull., U.S. 71(2):387-400.
- Bennett, D. H., J. A. Chandler, and L. K. Dunsmoor. 1990. Lower Granite Reservoir in-water disposal test: Results of the fishery, benthic, and habitat monitoring program - Year 1 (1988). Completion report to the U.S. Army Corps of Engineers, Walla Walla District, 251 p. (Available from Department of Fish and Wildlife Resources, College of Forestry, Wildlife and Range Sciences, University of Idaho, Moscow, Idaho 83843.)

Bennett, D. H., T. J. Dresser, T. S. Curet, K. B. Lepra, and M. A. Madsen. 1993. Lower Granite Reservoir in-water disposal test: Results of the fishery, benthic and habitat monitoring program - Year 4 (1991). Completion report to the U.S. Army Corps of Engineers, Walla Walla District, 234 p. (Available from Department of Fish and Wildlife Resources, College of Forestry, Wildlife and Range Sciences, University of Idaho, Moscow, Idaho 83843.)

Bennett, D. H., T. S. Curet, and M. A. Madsen. In prep. Abundance, habitat and migration of age-0 chinook salmon in the lower Snake River reservoirs with emphasis on Little Goose Reservoir, Washington. Report to the U.S. Army Corps of Engineers, Walla Walla District, 28 p. (Available from Department of Fish and Wildlife Resources, College of Forestry, Wildlife and Range Sciences, University of Idaho, Moscow, Idaho 83843.)

Bennett, D. H., T. L. Nightengale, and M. A. Madsen. 1997. Comparison and dynamics of benthic macroinvertebrate communities of Lower Granite, Little Goose and Lower Monumental reservoirs. Completion report to the U.S. Army Corps of Engineers, Walla Walla District. (Available from Department of Fish and Wildlife Resources, College of Forestry, Wildlife and Range Sciences, University of Idaho, Moscow, Idaho 83843.)

- Borror, D. J., D. M. DeLong, and C. A. Triplehorn. 1976. An introduction to the study of insects, fourth edition. Holt, Rinehart and Winston. U.S.A., 852 p.
- Bousfield, E. L. 1979. The amphipod superfamily Gammaroidea in the northeastern Pacific region: Systematics and distributional ecology. *Bull. Biol. Soc. Wash.* 3:297-357.
- Brinkhurst, R. O., and D. G. Cook. 1974. Aquatic earthworms (Annelida: Oligochaeta). *In* C. W. Hart, Jr. and S. L. H. Fuller (editors). *Pollution ecology of freshwater invertebrates*, p. 143-156. Academic Press, Inc. New York, NY.
- Brinkhurst, R. O., and S. R. Gelder. 1991. Annelida: Oligochaeta and Branchiobdellida. *In* J. H. Thorp and A. P. Covich (editors). *Ecology and classification of North American freshwater invertebrates*, p. 401-435. Academic Press, Inc. San Diego, CA.
- Dorband, W. R. 1980. Benthic macroinvertebrate communities in the lower Snake River reservoir system. Ph.D. Dissertation. University of Idaho, Moscow, ID, 160 p.
- Juul, S. T. J. 1997. Water quality update on the Lower Snake River, Washington. State of Washington Water Research Center, Report number 96. Washington State University, Pullman, Washington.

- Kirn, R. A., R. D. Ledgerwood, and A. L. Jensen. 1986. Diet of subyearling chinook salmon (*Oncorhynchus tshawytscha*) in the Columbia River estuary and changes effected by the 1980 eruption of Mount St. Helens. Northwest Sci. 60(3):191-196.
- Ledgerwood, R. D., S. S. Pool, and S. J. Grabowski. 1996. Limnological investigations in selected shallow-water habitats of Lower Granite Reservoir, 1995. Progress report to the U.S. Army Corps of Engineers, Walla Walla District, 43 p. (Available from Northwest Fisheries Science Center, 2725 Montlake Blvd. E., Seattle, WA 98112-2097.)
- Macdonald, J. S., and R. H. Green. 1983. Redundancy of variables used to describe importance of prey species in fish diets. Can. J. Fish. Aquat. Sci. 40(5):635-637.
- Muir, W. D., and R. L. Emmett. 1988. Food habits of migrating salmonid smolts passing Bonneville Dam in the Columbia River, 1984. Regul. Rivers Res. & Manage. 2:1-10.
- Muir, W. D. and T. C. Coley. 1996. Diet of yearling chinook salmon and feeding success during downstream migration in the Snake and Columbia Rivers. Northwest Sci. 70(4):298-305.
- Pennak, R. W. 1978. Fresh-water invertebrates of the United States, second edition. John Wiley & Sons, Inc. New York, NY, 803 p.

Roback, S. S. 1974. Insects (Arthropoda: Insecta). *In* C. W. Hart, Jr. and S. L. H. Fuller (editors). Pollution ecology of freshwater invertebrates, p. 314-376. Academic Press, Inc. New York, NY.

Rudy, P., Jr. and L. H. Rudy. 1983. Oregon estuarine invertebrates: An illustrated guide to the common and important invertebrate animals. U.S. Fish Wildl. Serv. Biol. Rep. Contract No. 79-111. FWS/OBS-83/16. 225 p.

Smith, R. I. and J. T. Carlton. 1975. Light's manual: Intertidal invertebrates of the central California coast, third edition. University of California Press, U.S.A., 716 p.

U.S. Army Corps of Engineers. 1992a. Lower Granite and Little Goose projects -- 1992 reservoir drawdown test report (draft, October 1992). Internal report to the U.S. Army Corps of Engineers, Walla Walla District, 138 p. (Available from U.S. Army Corps of Engineers, Walla Walla District, 201 North 3rd, Walla Walla, WA 99362-1876.)

U.S. Army Corps of Engineers. 1992b. Lower Granite Lock and Dam sedimentation removal for flood control preliminary evaluation and progress report (draft, December 1992). Internal report to the U.S. Army Corps of Engineers, Walla Walla District, 235 p. (Available from U.S. Army Corps of Engineers, Walla Walla District, 201 North 3rd, Walla Walla, WA 99362-1876.)

Word, J. Q. 1976. An evaluation of benthic invertebrate sampling devices for investigating feeding habits of fish. *In* C. A. Simenstad and S. J. Lipovsky (editors). Fish food habits studies, p. 43-55. WSG-WO 77-2. Washington Sea Grant, Division of Marine Resources, University of Washington, HG-30, Seattle, Washington.

APPENDICES

Appendix A

Subsampling procedure

Appendix B

**Habitat preferences of organisms
Crustacean densities
Sediment types and characteristic data**

Appendix C

Species compositions and densities

Appendix D

**Data analysis sheet
Spatial distribution**

APPENDICES

Appendix A


Appendix A1. Methodology used for subsampling large volume benthic invertebrate samples containing fine sediments.

A. Description

1. We used a modified Motoda plankton splitter to divide a large benthic invertebrate sample into six equivalent subsamples. Samples chosen for splitting had volumes about 1 to 2 L after sieving. Two or six subsamples were chosen for analysis. Each subsample was processed as an independent sample using standard procedures. In addition to the standard procedures, the subsample number and a comment clarifying that the sample processed was a subsample were included on the data sheet.
2. After each subsample was processed, tallies from each taxon/category were added then mathematically adjusted to estimate total tallies for the entire sample. The adjusted tallies were recorded on an independent data sheet, with the subsample data sheets attached.

B. Equipment

1. Motoda plankton splitter with 4-L capacity, 6 sub-divisions, 6 - 31.8 mm diameter rubber stoppers, and plastic vent covers.
2. U.S.A. Standard Testing Sieve no. 35 (0.5 mm)
3. Spoon
4. Wash bottle with tap water (500 ml)
5. Six jars with caps (250 to 500 ml)
6. Waterproof labels
7. Pencil
8. Scissors
9. Ethanol solution (80%)
10. Sink



Split compartment numbers

C. Rinse

1. Rinse the sample with tap water over a U.S.A. Standard Testing Sieve no. 35 (0.5 mm) to remove formalin.
2. Transfer the washed sample from the sieve to a holding jar. Add sufficient ethanol to cover the sediment.
3. Place the base of the plankton splitter on the counter so that the vents hang over the edge of the sink. Make the base of the splitter level using the corner screws and the lever indicator on the base.

4. Close each vent on the splitter with rubber stoppers and cover each stopper with a plastic vent cover to further seal the drains. Tilt the non-divided end of the splitter down on its rest.
5. Transfer the sample from the holding jar into the non-divided end of the splitter using the spoon.

D. Mixing and Splitting

1. Add sufficient tap water to the sample in the splitter to create a homogeneous slurry. Mix the slurry by rocking the splitter up and down several times.
2. Conclude mixing with the divided end of the splitter tilted down on its rest and the six vents extending over the edge of the sink. Check to see that each division has an equivalent amount of slurry, and if not, mix again.
3. Using the wash bottle with tap water, rinse the remaining sediment from the non-divided end of the splitter into the divided end. Use a side to side motion during this rinse to move the sediment into the divisions equally.

E. Subsample Acquisition

1. Hold the 0.5 mm sieve immediately under side vent number 1. Remove the stopper and plastic vent cover to let the subsample drain out onto the sieve. Restopper the vent, rinse completely, and drain again to remove all the sediment from the subcompartment.
2. Transfer the sample from the sieve to an appropriate sized jar. Add sufficient ethanol to the sample jar to cover the sediment. Place two waterproof paper labels with the original sample data and subsample number into the sample jar, and cap tightly.
3. The remaining subsamples are obtained from the other subcompartments in a similar manner.

Appendix B

Appendix Table B1. Non-insect taxa recovered from benthic invertebrate samples collected in Lower Granite Reservoir, 1994-1995. For data analysis purposes, the habitat type is noted for each organism (Barnard 1969, Smith and Carlton 1975, Borror et al. 1976, Pennak 1978, Bousfield 1979, Rudy and Rudy 1983, Barnes 1987).

Non-insect taxon	Benthic	Epibenthic	Planktonic	Terrestrial	Comments
Araneae				x	
Archaeogastropoda	x				
Bivalvia	x				
Calanoida			x		Mostly planktonic (Barnes 1987).
Cladocera	x		x		Was not consistently enumerated, therefore excluded from data analysis. It is mainly planktonic.
Copepoda	x		x		Habitat types vary between orders (Barnes 1987).
<i>Corophium salmonis</i>	x	x			Associates with muddy substrate (Rudy and Rudy 1983).
<i>Corophium spinicorne</i>	x	x			Associates with sandy substrate (Rudy and Rudy 1983).
<i>Corophium</i> spp.	x	x			Associates with mud and sand (Rudy and Rudy 1983).
Cyclopoida		x	x		Associates with substrate but also swims (Pennak 1978).
Eggs (unidentified)					Excluded from data analysis.
Gammaridae	x	x			Associates with substrate (Pennak 1978).
Gastropoda	x				
Harpacticoida	x				Crawls or runs on substrate and is restricted to bottom debris (Pennak 1978).
Hirudinea		x			Adheres to substrate for protection and for resting (Pennak 1978).
<i>Hyaella azteca</i>	x	x			Associates with substrate (Pennak 1978).
Isopoda	x				

Appendix Table B1. Continued.

Non-insect taxon	Benthic	Epibenthic	Planktonic	Terrestrial	Comments
Ixodides				x	
<i>Leptodora kindtii</i>			x		Excluded from data analysis.
Leptodoridae			x		Excluded from data analysis.
Mysidacea	x		x		Both benthic and planktonic forms exist (Barnes 1987).
Nematoda	x				Was not consistently enumerated, therefore excluded from data analysis.
Nematomorpha		x			Nematomorpha was misidentified as Nematoda, so it was excluded from analysis.
Nemertea	x				
Oligochaeta	x				
Ostracoda	x				
Polychaeta	x				
<i>Porcellio</i> spp.				x	
Prostigmata	x				
<i>Ramellogammarus oregonensis</i>	x	x			
<i>Ramellogammarus ramellus</i>	x	x			
Turbellaria		x			

Appendix Table B2. Insects recovered from benthic invertebrate samples collected in Lower Granite Reservoir, 1994-1995. For data analysis purposes, the life stages of each insect taxon is noted as to whether it is benthic (Borror et al. 1976, Pennak 1978, Barnes 1987).

Taxon	Larva/ nymph	Pupa	Adult	Comments
Caenidae	x	n/a*		
<i>Caenis</i> spp.	x	n/a		
Ceratopogonidae	x	x		The aquatic immature stages are generally benthic.
Chironomidae	x	x		The aquatic immature stages are generally benthic.
Chironominae	x	x		The aquatic immature stages are generally benthic.
Coleoptera	x		x	
Collembola				Some species are aquatic and restricted to the water surface. No reference to benthic species were found.
Culicidae				The larval form generally stays at the water surface. Hence, no life stage was included as benthic.
Diptera	x	x		
Elmidae	x		x	
<i>Ephemera</i> spp.	x	n/a		
Ephemeroptera	x	n/a		
Formicidae				
Hemiptera				Most species are terrestrial and some are aquatic.
<i>Hexagenia</i> spp.	x	n/a		
Homoptera		n/a		
Insecta	x	x	x	Organisms recorded as Insecta were not further identified due to fragmentation, therefore all of these were marked as benthic invertebrates.
Isoptera				
Lepidoptera larvae	x			
Leptophlebiidae	x	n/a		
Orthoclaadiinae	x	x		The aquatic immature stages are generally benthic.
Orthoptera	x	x		
Plecoptera	x	n/a		
Psocoptera		n/a		
Psychomyiidae	x	x		

Appendix Table B2. Continued

Taxon	Larva/ nymph	Pupa	Adult	Comments
<i>Sialis</i> spp.	x			<i>Sialis</i> spp. can be associated with substrate (Pennak 1978).
Simuliidae				The immature life stages are found in shallow water of mostly swift streams (Pennak 1978).
Tanyderidae larvae	x			
Tanypodinae	x	x		The aquatic immature stages are generally benthic.
Thysanoptera				
Trichoptera	x	x		

^a The "n/a" in the pupa column denotes insects having a nymphal, not pupal life stage.

Appendix Table B3. Mean densities (no./m²) of Crustacea taxa recovered from the benthic invertebrate samples collected in Lower Granite Reservoir, 1994-1995.

Sampling area	Depth (m)	Date	Amphipoda	Copepoda	Isopoda	Mysidacea	Ostracoda
Silcott Island	3	30 Mar 94	0	25	0	0	5
		4 May 94	0	135	0	0	0
		1 Jun 94	10	0	0	0	0
		29 Jun 94	0	275	0	0	10
		27 Jul 94	0	5	0	0	0
		24 Aug 94	0	40	0	0	35
		21 Sep 94	0	50	5	0	15
		26 Oct 94	0	30	0	0	15
		30 Nov 94	5	60	0	0	70
		18 Jan 95	0	255	0	0	30
		22 Mar 95	0	25	0	0	50
		26 Apr 95	0	65	0	0	0
		24 May 95	0	30	0	0	0
		21 Jun 95	45	15	0	0	45
		19 Jul 95	0	0	0	0	80
		16 Aug 95	0	210	0	0	15
		13 Sep 95	0	15	0	0	10
		11 Oct 95	0	50	0	0	35
		Mean:	3	71	<1	0	23
Centennial Island	3	30 Mar 94	5	0	0	0	15
		4 May 94	0	5	0	0	145
		1 Jun 94	15	0	0	0	10
		29 Jun 94	0	10	0	0	0
		27 Jul 94	0	0	0	0	5
		24 Aug 94	0	185	0	0	85
		21 Sep 94	0	0	0	0	30
		26 Oct 94	0	0	0	0	0
		30 Nov 94	0	295	0	0	25
		18 Jan 95	5	0	0	0	30
		22 Mar 95	0	0	0	0	55
		26 Apr 95	0	10	0	0	60
		24 May 95	10	105	0	0	25
		21 Jun 95	0	5	0	0	0
		19 Jul 95	0	15	0	0	25
		16 Aug 95	0	50	0	0	50
		13 Sep 95	10	5	0	0	40
		11 Oct 95	0	0	0	0	10
		Mean:	3	38	0	0	34
Offfield	3	30 Mar 94	10	10	0	0	0
		4 May 94	0	30	0	0	100
		1 Jun 94	5	65	0	0	5
		29 Jun 94	5	100	10	0	5
		27 Jul 94	0	10	0	0	25
		24 Aug 94	10	10	10	0	30
		21 Sep 94	0	25	0	0	10
		26 Oct 94	0	0	0	0	5
		30 Nov 94	0	5	0	0	30

Appendix Table B3. Continued.

Sampling area	Depth (m)	Date	Amphipoda	Copepoda	Isopoda	Mysidacea	Ostracoda
		18 Jan 95	0	10	0	0	30
		22 Mar 95	5	0	0	0	10
		26 Apr 95	620	40	0	0	35
		24 May 95	0	0	0	0	15
		21 Jun 95	0	0	0	0	10
		19 Jul 95	0	65	0	0	30
		16 Aug 95	10	70	0	0	15
		13 Sep 95	10	0	0	0	10
		11 Oct 95	20	0	0	0	5
		Mean:	39	24	1	0	21
Silcott Island	9	30 Mar 94	0	0	0	0	0
		4 May 94	0	0	0	0	0
		1 Jun 94	0	25	0	0	5
		29 Jun 94	0	20	0	0	5
		27 Jul 94	0	50	0	0	0
		24 Aug 94	10	115	0	0	35
		21 Sep 94	5	25	0	0	115
		26 Oct 94	5	0	0	0	5
		30 Nov 94	5	20	0	0	90
		18 Jan 95	0	50	0	0	45
		22 Mar 95	0	5	0	0	0
		26 Apr 95	0	45	0	0	5
		24 May 95	0	30	0	0	0
		21 Jun 95	0	0	0	0	0
		19 Jul 95	0	10	0	0	135
		16 Aug 95	0	25	0	0	25
		13 Sep 95	0	0	0	0	15
		11 Oct 95	0	25	0	0	0
		Mean:	1	25	0	0	27
Centennial Island	9	30 Mar 94	10	25	0	0	15
		4 May 94	5	0	0	0	25
		1 Jun 94	0	205	0	0	40
		29 Jun 94	0	85	0	0	0
		27 Jul 94	0	0	0	0	0
		24 Aug 94	0	180	0	0	50
		21 Sep 94	0	10	0	25	0
		26 Oct 94	5	0	0	0	5
		30 Nov 94	0	100	0	0	30
		18 Jan 95	10	0	0	0	10
		22 Mar 95	5	0	0	0	10
		26 Apr 95	0	355	0	0	50
		24 May 95	0	390	0	0	25
		21 Jun 95	5	195	0	0	0
		19 Jul 95	0	50	0	0	25
		16 Aug 95	0	25	0	0	25
		13 Sep 95	0	15	0	0	10
		11 Oct 95	0	15	0	0	0
		Mean:	2	92	0	1	18

Appendix Table B3. Continued.

Sampling area	Depth (m)	Date	Amphipoda	Copepoda	Isopoda	Mysidacea	Ostracoda
Offfield	9	30 Mar 94	5	15	0	0	0
		4 May 94	0	0	0	0	0
		1 Jun 94	0	10	0	0	5
		29 Jun 94	5	65	0	0	0
		27 Jul 94	0	0	10	0	0
		24 Aug 94	5	0	0	0	55
		21 Sep 94	15	0	0	0	50
		26 Oct 94	15	0	0	0	0
		30 Nov 94	195	275	0	0	365
		18 Jan 95	120	0	0	0	115
		22 Mar 95	5	0	0	0	5
		26 Apr 95	430	25	0	0	25
		24 May 95	25	50	0	0	25
		21 Jun 95	0	0	0	0	20
		19 Jul 95	10	40	0	0	10
		16 Aug 95	10	5	0	0	15
		13 Sep 95	0	5	0	0	0
		11 Oct 95	0	5	0	0	10
		Mean:	47	28	1	0	39
Silcott Island	18	30 Mar 94	5	0	0	0	0
		4 May 94	5	0	0	0	0
		1 Jun 94	0	0	0	0	0
		29 Jun 94	0	0	0	0	0
		27 Jul 94	0	0	0	0	35
		24 Aug 94	0	230	0	0	125
		21 Sep 94	75	0	0	0	150
		26 Oct 94	0	0	0	0	0
		30 Nov 94	0	15	0	0	140
		18 Jan 95	0	0	0	0	90
		22 Mar 95	0	0	0	0	10
		26 Apr 95	0	0	0	0	45
		24 May 95	0	0	0	0	15
		21 Jun 95	0	0	0	0	0
		19 Jul 95	0	5	0	0	0
		16 Aug 95	5	0	0	0	0
		13 Sep 95	0	0	0	0	0
		11 Oct 95	0	5	0	0	0
		Mean:	5	14	0	0	34
Centennial Island	18	30 Mar 94	0	210	0	0	10
		4 May 94	0	0	0	0	5
		1 Jun 94	0	20	0	0	5
		29 Jun 94	0	105	0	0	15
		27 Jul 94	0	0	0	0	0
		24 Aug 94	0	370	0	0	25
		21 Sep 94	0	0	0	0	15
		26 Oct 94	0	5	0	0	0
		30 Nov 94	0	35	0	0	35
		18 Jan 95	0	0	0	0	10

Appendix Table B3. Continued.

Sampling area	Depth (m)	Date	Amphipoda	Copepoda	Isopoda	Mysidacea	Ostracoda
		22 Mar 95	0	0	0	0	0
		26 Apr 95	0	10	0	0	170
		24 May 95	0	175	0	0	40
		21 Jun 95	0	20	0	0	5
		19 Jul 95	0	15	0	0	20
		16 Aug 95	5	0	0	0	5
		13 Sep 95	0	0	0	0	5
		11 Oct 95	0	0	0	0	0
		Mean:	<1	54	0	0	20
Offfield	18	30 Mar 94	0	25	0	0	40
		4 May 94	0	0	0	0	5
		1 Jun 94	0	0	0	0	0
		29 Jun 94	0	5	0	0	20
		27 Jul 94	180	15	0	0	80
		24 Aug 94	5	65	0	0	230
		21 Sep 94	15	0	0	0	185
		26 Oct 94	55	0	0	0	10
		30 Nov 94	45	310	0	0	70
		18 Jan 95	140	175	0	0	50
		22 Mar 95	45	70	0	0	35
		26 Apr 95	15	50	0	0	65
		24 May 95	0	5	0	0	65
		21 Jun 95	0	0	0	0	15
		19 Jul 95	40	5	0	0	5
		16 Aug 95	0	225	0	0	0
		13 Sep 95	0	10	0	0	15
		11 Oct 95	5	5	0	0	15
		Mean:	30	54	0	0	50

Appendix Table B4. Percentages of gravel, sand, and fines in sediments collected in July 1995 from all benthic invertebrate sampling stations in Lower Granite Reservoir.

Sampling area	Depth (m)	Sediment type ^b	Benthic invertebrate transect ^a			
			Upstream	Middle upstream	Middle downstream	Downstream
Silcott Island (RKm 212)	3	Gravel (%)	0.0	0.0	0.0	0.0
		Sand (%)	15.7	40.0	25.8	12.9
		Fines (%)	84.3	60.0	74.2	87.1
	9	Gravel (%)	0.0	0.0	0.0	0.0
		Sand (%)	61.1	24.2	55.9	17.8
		Fines (%)	38.9	75.8	44.1	82.2
	18	Gravel (%)	0.0	0.0	0.0	0.0
		Sand (%)	94.0	93.9	91.5	37.3
		Fines (%)	6.0	6.1	8.5	62.7
Centennial Island (RKm 193)	3	Gravel (%)	0.0	0.0	0.0	0.0
		Sand (%)	72.4	79.6	72.5	71.1
		Fines (%)	27.6	20.4	27.5	28.9
	9	Gravel (%)	0.0	0.0	0.0	0.0
		Sand (%)	4.3	6.9	61.6	10.6
		Fines (%)	95.7	93.1	38.4	89.4
	18	Gravel (%)	0.0	0.0	0.0	0.0
		Sand (%)	5.4	6.7	6.7	7.5
		Fines (%)	94.5	93.3	93.3	92.5
Offfield (RKm 177)	3	Gravel (%)	3.7	0.0	2.2	7.4
		Sand (%)	28.6	19.1	31.0	58.0
		Fines (%)	67.6	80.9	66.8	34.7
	9	Gravel (%)	2.6	0.0	0.0	38.8
		Sand (%)	42.9	8.3	6.2	19.2
		Fines (%)	54.6	91.7	93.8	42.0
	18	Gravel (%)	3.0	0.0	28.5	0.0
		Sand (%)	13.0	32.4	8.5	7.1
		Fines (%)	84.0	67.6	63.0	92.9

^a Refer to Figure 2 in main text for details.

^b Analysis under contract by the U.S. Army Corps of Engineers North Pacific Division Materials Laboratory, Troutdale, Oregon.

Appendix Table B5. Median grain sizes (mm), silt/clay (%), and volatile solids (%) in sediments collected in July 1995 from benthic invertebrate sampling stations in Lower Granite Reservoir.

Sampling area	Depth (m)	Sediment characteristic ^b	Benthic invertebrate transect ^a			
			Upstream	Middle upstream	Middle downstream	Downstream
Silcott Island (RKm 212)	3	Median grain size (mm)	0.06	0.07	0.06	0.03
		Silt/clay (%)	82.0	47.0	63.1	84.5
		Volatile solids (%)	6.6	6.2	6.4	17.8
	9	Median grain size (mm)	0.09	0.06	0.08	0.04
		Silt/clay (%)	27.4	73.0	28.8	74.6
		Volatile solids (%)	3.5	3.4	3.1	7.4
	18	Median grain size (mm)	0.18	0.18	0.18	0.05
		Silt/clay (%)	5.4	5.4	8.4	58.6
		Volatile solids (%)	0.4	0.7	0.7	48.3
Centennial Island (RKm 193)	3	Median grain size (mm)	0.26	0.21	0.28	0.14
		Silt/clay (%)	27.5	20.3	26.8	24.9
		Volatile solids (%)	3.4	2.1	3.8	3.4
	9	Median grain size (mm)	0.02	0.02	0.14	0.02
		Silt/clay (%)	95.0	91.0	37.0	86.8
		Volatile solids (%)	13.4	15.8	4.2	11.6
	18	Median grain size (mm)	0.02	0.02	0.02	0.02
		Silt/clay (%)	93.5	90.9	91.1	90.4
		Volatile solids (%)	12.1	11.3	10.9	13.0
Offfield (RKm 177)	3	Median grain size (mm)	0.05	0.05	0.05	0.15
		Silt/clay (%)	59.9	77.5	59.7	32.4
		Volatile solids (%)	5.2	4.9	3.8	5.1
	9	Median grain size (mm)	0.06	0.04	0.03	0.36
		Silt/clay (%)	52.1	90.6	93.5	41.3
		Volatile solids (%)	6.2	6.1	6.6	6.8
	18	Median grain size (mm)	0.03	0.04	0.03	0.01
		Silt/clay (%)	79.5	62.3	61.6	90.6
		Volatile solids (%)	7.3	6.8	9.6	10.0

^a Refer to Figure 2 in main text for details.

^b Analysis under contract by the U.S. Army Corps of Engineers North Pacific Division Materials Laboratory, Troutdale, Oregon.

Appendix C

Appendix Table C1. Species compositions and densities (no./m²) by date, depth, and sampling area in Lower Granite Reservoir, 1994-1995.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
Silcott Island				4 May 94			
30 Mar 94				9 m	No. samples processed:	4	
3 m	No. samples processed:	4		Oligochaeta		455	2,275
Oligochaeta		3,261	16,305	Chironomidae larvae		70	350
Chironomidae larvae		297	1,485	Cladocera		6	30
Nematoda		24	120	Chironomidae pupae		4	20
Cyclopoida		14	70	Trichoptera larvae		3	15
Copepoda		5	25	Bivalvia		2	10
Prostigmata (water mites)		4	20	Diptera adult		2	10
Bivalvia		3	15	Nematoda		1	5
Nemertea		2	10	Nematomorpha		1	5
Ephemeroptera nymph		1	5	Nemertea		1	5
Ostracoda		1	5	Polychaeta		1	5
Turbellaria		1	5	Prostigmata (water mites)		1	5
Subtotal:		3,613	18,065	Subtotal:		547	2,735
30 Mar 94				4 May 94			
9 m	No. samples processed:	3		18 m	No. samples processed:	4	
Oligochaeta		1,018	6,787	Oligochaeta		468	2,340
Chironomidae larvae		174	1,160	Eggs (unidentified)		17	85
Nematoda		4	27	Chironomidae larvae		8	40
Cyclopoida		1	7	<i>Corophium salmonis</i>		1	5
Ephemeroptera nymph		1	7	Trichoptera larvae		1	5
Subtotal:		1,198	7,987	Subtotal:		495	2,475
30 Mar 94				Subtotal all depths for this sampling area and date period:			
18 m	No. samples processed:	4				2,686	4,477
Oligochaeta		931	4,655				
Chironomidae larvae		81	405	1 Jun 94			
Cladocera		2	10	3 m	No. samples processed:	4	
Turbellaria		2	10	Oligochaeta		1,275	6,375
Araneae (spiders)		1	5	Chironomidae larvae		235	1,175
<i>Corophium spinicorne</i>		1	5	Nematoda		16	80
Subtotal:		1,018	5,090	Cladocera		7	35
Subtotal all depths for this sampling area and date period:		5,829	10,381	Prostigmata (water mites)		5	25
4 May 94				Ephemeroptera nymph		4	20
3 m	No. samples processed:	4		Collembola		3	15
Oligochaeta		1,406	7,030	Nemertea		3	15
Chironomidae larvae		101	505	Simuliidae larvae		3	15
Cladocera		39	195	<i>Porcellio</i> spp.		2	10
Nematoda		29	145	Archaeogastropoda		1	5
Copepoda		27	135	Bivalvia		1	5
Eggs (unidentified)		27	135	Chironomidae pupae		1	5
Nemertea		8	40	<i>Corophium salmonis</i>		1	5
Insecta		3	15	<i>Corophium</i> spp.		1	5
Turbellaria		3	15	Cyclopoida		1	5
Diptera pupae		1	5	Plecoptera nymph		1	5
Subtotal:		1,644	8,220	Trichoptera larvae		1	5
				Subtotal:		1,561	7,805

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
1 Jun 94 9 m	No. samples processed:	4			Cyclopoida	4	20
	Oligochaeta	789	3,945		Copepoda	3	15
	Chironomidae larvae	196	980		Chironomidae pupae	2	10
	Nematoda	22	110		Polychaeta	2	10
	Harpacticoida	5	25		Bivalvia	1	5
	Ephemeroptera nymph	4	20		Cladocera	1	5
	Coleoptera larvae	3	15		Coleoptera adult	1	5
	Cyclopoida	3	15		Harpacticoida	1	5
	Prostigmata (water mites)	2	10		Ostracoda	1	5
	Bivalvia	1	5		Subtotal:	1,889	9,445
	Chironomidae pupae	1	5	29 Jun 94 18 m	No. samples processed:	4	
	Formicidae adult	1	5		Oligochaeta	421	2,105
	Ostracoda	1	5		Chironomidae larvae	34	170
	Trichoptera larvae	1	5		Eggs (unidentified)	11	55
	Subtotal:	1,029	5,145		Bivalvia	7	35
1 Jun 94 18 m	No. samples processed:	4			Cladocera	3	15
	Oligochaeta	146	730		Polychaeta	3	15
	Chironomidae larvae	5	25		Unidentified	3	15
	Eggs (unidentified)	1	5		Diptera pupae	1	5
	Nematoda	1	5		Nemertea	1	5
	Plecoptera nymph	1	5		Subtotal:	484	2,420
	Subtotal:	154	770		Subtotal all depths for this sampling area and date period:	6,428	10,713
	Subtotal all depths for this sampling area and date period:	2,744	4,573	27 Jul 94 3 m	No. samples processed:	4	
29 Jun 94 3 m	No. samples processed:	4			Oligochaeta	545	2,725
	Oligochaeta	3,811	19,055		Chironomidae larvae	37	185
	Cladocera	99	495		Chironomidae pupae	5	25
	Chironomidae larvae	50	250		Cyclopoida	5	25
	Harpacticoida	40	200		Nematoda	4	20
	Cyclopoida	21	105		Harpacticoida	1	5
	Copepoda	15	75		Subtotal:	597	2,985
	Turbellaria	5	25	27 Jul 94 9 m	No. samples processed:	4	
	Nematoda	3	15		Oligochaeta	1,484	7,420
	Nemertea	3	15		Chironomidae larvae	56	280
	Ostracoda	2	10		Cyclopoida	16	80
	Prostigmata (water mites)	2	10		Copepoda	9	45
	Chironomidae pupae	1	5		Chironomidae pupae	3	15
	Collembola	1	5		Hirudinea	2	10
	Ixodides (ticks)	1	5		Nematoda	2	10
	Trichoptera larvae	1	5		Bivalvia	1	5
	Subtotal:	4,055	20,275		Harpacticoida	1	5
29 Jun 94 9 m	No. samples processed:	4			Plecoptera nymph	1	5
	Oligochaeta	1,778	8,890		Polychaeta	1	5
	Nemertea	54	270		Subtotal:	1,576	7,880
	Chironomidae larvae	41	205	27 Jul 94 18 m	No. samples processed:	4	

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
	Oligochaeta	604	3,020	24 Aug 94			
	Chironomidae larvae	70	350	18 m	No. samples processed:	4	
	Ostracoda	7	35		Oligochaeta	693	3,465
	Cyclopoida	5	25		Chironomidae larvae	100	500
	Chironomidae pupae	2	10		Copepoda	46	230
	Bivalvia	1	5		Ostracoda	25	125
	Ephemeroptera nymph	1	5		Cyclopoida	18	90
	Nematoda	1	5		Eggs (unidentified)	13	65
	Polychaeta	1	5		Cladocera	6	30
	Prostigmata (water mites)	1	5		Bivalvia	5	25
	Subtotal:	693	3,465		Diptera pupae	2	10
	Subtotal all depths for this sampling area and date period:	2,866	4,777		Ceratopogonidae larvae	1	5
24 Aug 94					Chironomidae pupae	1	5
3 m	No. samples processed:	4			Diptera larvae	1	5
	Oligochaeta	1,659	8,295		Hirudinea	1	5
	Chironomidae larvae	632	3,160		Polychaeta	1	5
	Cladocera	81	405		Subtotal:	913	4,565
	Eggs (unidentified)	37	185		Subtotal all depths for this sampling area and date period:	6,590	10,983
	<i>Leptodora kindtii</i>	13	65	21 Sep 94			
	Copepoda	7	35	3 m	No. samples processed:	4	
	Ostracoda	7	35		Oligochaeta	3,708	18,540
	Chironomidae pupae	5	25		Chironomidae larvae	295	1,475
	Cyclopoida	3	15		Harpacticoida	9	45
	Nematoda	3	15		Cyclopoida	5	25
	Diptera pupae	2	10		Nematoda	5	25
	Bivalvia	1	5		Ostracoda	3	15
	Chironomidae adult	1	5		Bivalvia	2	10
	Harpacticoida	1	5		Chironomidae pupae	1	5
	Subtotal:	2,452	12,260		Copepoda	1	5
24 Aug 94					Isopoda	1	5
9 m	No. samples processed:	4			Subtotal:	4,030	20,150
	Oligochaeta	1,346	6,730	21 Sep 94			
	Cladocera	1,094	5,470	9 m	No. samples processed:	4	
	Chironomidae larvae	641	3,205		Oligochaeta	4,854	24,270
	Cyclopoida	65	325		Chironomidae larvae	402	2,010
	<i>Leptodora kindtii</i>	34	170		Ostracoda	23	115
	Copepoda	20	100		Nematoda	10	50
	Ostracoda	7	35		Chironomidae pupae	5	25
	Chironomidae pupae	4	20		Cyclopoida	5	25
	Harpacticoida	3	15		Bivalvia	4	20
	Nematoda	3	15		Harpacticoida	4	20
	Bivalvia	2	10		Araneae (spiders)	1	5
	Polychaeta	2	10		Copepoda	1	5
	<i>Corophium spinicorne</i>	1	5		<i>Corophium salmonis</i>	1	5
	Orthocladinae pupae	1	5		Prostigmata (water mites)	1	5
	<i>Ramullogammarus oregonensis</i>	1	5		Subtotal:	5,311	26,555
	Turbellaria	1	5	21 Sep 94			
	Subtotal:	3,225	16,125	18 m	No. samples processed:	4	

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
	Oligochaeta	789	3,945		Oligochaeta	5,725	28,625
	Chironomidae larvae	188	940		Chironomidae larvae	310	1,550
	Ostracoda	30	150		Cyclopoida	29	145
	<i>Corophium spinicorne</i>	14	70		Nematoda	14	70
	Polychaeta	13	65		Ostracoda	14	70
	Bivalvia	10	50		Copepoda	6	30
	Cyclopoida	7	35		Harpacticoida	6	30
	Nemertea	4	20		Archaeogastropoda	3	15
	Chironomidae pupae	2	10		Turbellaria	2	10
	Cladocera	1	5		Araneae (spiders)	1	5
	<i>Corophium salmonis</i>	1	5		Bivalvia	1	5
	Subtotal:	1,059	5,295		Calanoida	1	5
	Subtotal all depths for this sampling area and date period:	10,400	17,333		Diptera	1	5
26 Oct 94					Ephemeroptera nymph	1	5
3 m	No. samples processed:	4			<i>Ramellogammarus oregonensis</i>	1	5
	Oligochaeta	2,263	11,315		Subtotal:	6,115	30,575
	Chironomidae larvae	138	690	30 Nov 94			
	Nematoda	7	35	9 m	No. samples processed:	4	
	Harpacticoida	6	30		Oligochaeta	3,044	15,220
	Ostracoda	3	15		Chironomidae larvae	244	1,220
	Bivalvia	2	10		Ostracoda	18	90
	Cyclopoida	2	10		Bivalvia	3	15
	Prostigmata (water mites)	2	10		Copepoda	3	15
	Araneae (spiders)	1	5		Cyclopoida	3	15
	Collembola	1	5		Nematoda	2	10
	Trichoptera larvae	1	5		Prostigmata (water mites)	2	10
	Turbellaria	1	5		<i>Corophium salmonis</i>	1	5
	Subtotal:	2,427	12,135		Harpacticoida	1	5
26 Oct 94					Subtotal:	3,321	16,605
9 m	No. samples processed:	4		30 Nov 94			
	Oligochaeta	1,093	5,465	18 m	No. samples processed:	4	
	Chironomidae larvae	163	815		Oligochaeta	1,017	5,085
	Bivalvia	3	15		Chironomidae larvae	259	1,295
	Nematoda	1	5		Ostracoda	28	140
	Ostracoda	1	5		Bivalvia	3	15
	<i>Ramellogammarus ramellus</i>	1	5		Copepoda	3	15
	Subtotal:	1,262	6,310		Eggs (unidentified)	3	15
26 Oct 94					Nematoda	3	15
18 m	No. samples processed:	4			Cyclopoida	2	10
	Oligochaeta	1,015	5,075		Nemertea	2	10
	Chironomidae larvae	81	405		Turbellaria	1	5
	Homoptera adult	1	5		Subtotal:	1,321	6,605
	Insecta adult	1	5		Subtotal all depths for this sampling area and date period:	10,757	17,928
	Subtotal:	1,098	5,490	18 Jan 95			
	Subtotal all depths for this sampling area and date period:	4,787	7,978	3 m	No. samples processed:	4	
30 Nov 94					Oligochaeta	6,647	33,235
3 m	No. samples processed:	4			Chironomidae larvae	336	1,680
					Turbellaria	201	1,005

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
	Nematoda	81	405		Harpacticoida	5	25
	Cyclopoida	63	315		Diptera pupae	3	15
	Copepoda	38	190		Trichoptera larvae	3	15
	Harpacticoida	13	65		Bivalvia	2	10
	Ostracoda	6	30		Hirudinea	1	5
	Bivalvia	3	15		Subtotal:	5,955	29,775
	Nemertea	3	15	22 Mar 95			
	Subtotal:	7,391	36,955	9 m	No. samples processed:	4	
18 Jan 95					Oligochaeta	2,997	14,985
9 m	No. samples processed:	4			Chironomidae larvae	215	1,075
	Oligochaeta	3,306	16,530		Nematoda	128	640
	Chironomidae larvae	253	1,265		Turbellaria	24	120
	Nematoda	43	215		Cyclopoida	7	35
	Cyclopoida	19	95		Prostigmata (water mites)	6	30
	Copepoda	10	50		Plecoptera nymph	3	15
	Ostracoda	9	45		Trichoptera larvae	3	15
	Nemertea	4	20		Ceratopogonidae larvae	1	5
	Bivalvia	3	15		Ephemeroptera nymph	1	5
	Hirudinea	3	15		Gastropoda	1	5
	Prostigmata (water mites)	3	15		Harpacticoida	1	5
	Araneae (spiders)	1	5		Nemertea	1	5
	Calanoida	1	5		Subtotal:	3,388	16,940
	Ephemeroptera nymph	1	5	22 Mar 95			
	Trichoptera larvae	1	5	18 m	No. samples processed:	4	
	Subtotal:	3,657	18,285		Oligochaeta	2,140	10,700
18 Jan 95					Chironomidae larvae	114	570
18 m	No. samples processed:	4			Nematoda	5	25
	Oligochaeta	3,099	15,495		Nemertea	5	25
	Chironomidae larvae	232	1,160		Ostracoda	2	10
	Ostracoda	18	90		Subtotal:	2,266	11,330
	Nematoda	7	35		Subtotal all depths for this sampling area and date period:	11,609	19,348
	Bivalvia	6	30	26 Apr 95			
	Cyclopoida	3	15	3 m	No. samples processed:	4	
	Nemertea	3	15		Oligochaeta	2,467	12,335
	Trichoptera larvae	1	5		Chironomidae larvae	249	1,245
	Subtotal:	3,369	16,845		Nematoda	46	230
	Subtotal all depths for this sampling area and date period:	14,417	24,028		Prostigmata (water mites)	11	55
22 Mar 95					Copepoda	10	50
3 m	No. samples processed:	4			Lepidoptera larvae	9	45
	Oligochaeta	5,354	26,770		Cyclopoida	6	30
	Chironomidae larvae	316	1,580		Coleoptera larvae	3	15
	Nematoda	111	555		Harpacticoida	3	15
	Turbellaria	72	360		Trichoptera larvae	2	10
	Cyclopoida	41	205		Bivalvia	1	5
	Nemertea	25	125		Hemiptera adult	1	5
	Ostracoda	10	50		Turbellaria	1	5
	Prostigmata (water mites)	7	35		Subtotal:	2,809	14,045
	Ephemeroptera nymph	5	25				

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
26 Apr 95							
9 m	No. samples processed:	4				Subtotal:	2,619 13,095
	Oligochaeta	2,645	13,225	24 May 95			
	Chironomidae larvae	273	1,365	18 m	No. samples processed:	4	
	Nematoda	28	140		Oligochaeta	378	1,890
	Nemertea	7	35		Chironomidae larvae	11	55
	Copepoda	6	30		Bivalvia	7	35
	Bivalvia	4	20		Ostracoda	3	15
	Prostigmata (water mites)	4	20		Chironomidae pupae	2	10
	Chironomidae pupae	3	15		Nematoda	2	10
	Harpacticoida	3	15		Ixodides (ticks)	1	5
	Collembola	1	5		Plecoptera nymph	1	5
	Ephemeroptera nymph	1	5		Subtotal:	405	2,025
	Ostracoda	1	5		Subtotal all depths for this sampling area and date period:	6,944	11,573
	Polychaeta	1	5	21 Jun 95			
	Subtotal:	2,977	14,885	3 m	No. samples processed:	4	
26 Apr 95					Oligochaeta	2,739	13,695
18 m	No. samples processed:	4			Chironomidae larvae	132	660
	Oligochaeta	1,767	8,835		Nematoda	78	390
	Chironomidae larvae	47	235		Bivalvia	51	255
	Bivalvia	9	45		Prostigmata (water mites)	42	210
	Ostracoda	9	45		Diptera pupae	15	75
	Nematoda	8	40		Cyclopoida	9	45
	Cyclopoida	2	10		Ephemeroptera nymph	9	45
	Nemertea	1	5		Ostracoda	9	45
	Prostigmata (water mites)	1	5		<i>Corophium</i> spp.	6	30
	Subtotal:	1,844	9,220		Hemiptera	6	30
	Subtotal all depths for this sampling area and date period:	7,630	12,717		Plecoptera nymph	6	30
24 May 95					Turbellaria	6	30
3 m	No. samples processed:	4			Archaeogastropoda	3	15
	Oligochaeta	3,689	18,445		Chironomidae adult	3	15
	Chironomidae larvae	174	870		<i>Corophium spinicorne</i>	3	15
	Nematoda	41	205		Diptera adult	3	15
	Copepoda	6	30		Harpacticoida	3	15
	Trichoptera larvae	6	30		Isoptera	3	15
	Chironomidae pupae	3	15		Lepidoptera larvae	3	15
	Ephemeroptera nymph	1	5		Trichoptera larvae	3	15
	Subtotal:	3,920	19,600		Subtotal:	3,132	15,660
24 May 95				21 Jun 95			
9 m	No. samples processed:	4		9 m	No. samples processed:	4	
	Oligochaeta	2,503	12,515		Oligochaeta	756	3,780
	Chironomidae larvae	93	465		Bivalvia	21	105
	Bivalvia	13	65		Chironomidae larvae	12	60
	Harpacticoida	6	30		Nematoda	5	25
	Ephemeroptera nymph	1	5		Trichoptera larvae	2	10
	Nematoda	1	5		Chironomidae pupae	1	5
	Prostigmata (water mites)	1	5		Coleoptera larvae	1	5
	Trichoptera larvae	1	5		Hemiptera	1	5
					Subtotal:	799	3,995

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
21 Jun 95 18 m	No. samples processed:	4		16 Aug 95 3 m	No. samples processed:	4	
	Oligochaeta	143	715		Oligochaeta	5,490	27,450
	Bivalvia	16	80		Chironomidae larvae	69	345
	Chironomidae larvae	2	10		Cyclopoida	47	235
	Diptera pupae	1	5		Copepoda	24	120
	Ephemeroptera nymph	1	5		Nematoda	19	95
	Subtotal:	163	815		Harpacticoida	18	90
	Subtotal all depths for this sampling area and date period:	4,094	6,823		Bivalvia	5	25
19 Jul 95 3 m	No. samples processed:	4			Ostracoda	3	15
	Oligochaeta	3,715	18,575		Tanyderidae larvae	3	15
	Nematoda	47	235		Turbellaria	3	15
	Chironomidae larvae	36	180		Subtotal:	5,681	28,405
	Ostracoda	16	80	16 Aug 95 9 m	No. samples processed:	4	
	Bivalvia	6	30		Oligochaeta	2,487	12,435
	Cyclopoida	6	30		Chironomidae larvae	45	225
	Hemiptera	6	30		Cyclopoida	24	120
	Prostigmata (water mites)	3	15		Plecoptera nymph	12	60
	Trichoptera larvae	3	15		Turbellaria	12	60
	Diptera	1	5		Copepoda	5	25
	Subtotal:	3,839	19,195		Ostracoda	5	25
19 Jul 95 9 m	No. samples processed:	4			Nematoda	3	15
	Oligochaeta	2,994	14,970		Chironomidae pupae	1	5
	Chironomidae larvae	72	360		Subtotal:	2,594	12,970
	Ostracoda	27	135	16 Aug 95 18 m	No. samples processed:	4	
	Polychaeta	8	40		Oligochaeta	299	1,495
	Bivalvia	2	10		Chironomidae larvae	29	145
	Copepoda	2	10		Bivalvia	10	50
	Ephemeroptera nymph	2	10		<i>Corophium salmonis</i>	1	5
	Nematoda	1	5		Nematoda	1	5
	Nemertea	1	5		No invertebrates present	0	0
	Prostigmata (water mites)	1	5		Subtotal:	340	1,700
	Subtotal:	3,110	15,550		Subtotal all depths for this sampling area and date period:	8,615	14,358
19 Jul 95 18 m	No. samples processed:	4		13 Sep 95 3 m	No. samples processed:	4	
	Oligochaeta	533	2,665		Oligochaeta	2,404	12,020
	Chironomidae larvae	37	185		Chironomidae larvae	154	770
	Polychaeta	12	60		Nematoda	5	25
	Bivalvia	9	45		Harpacticoida	3	15
	Unidentified	7	35		Nemertea	3	15
	Chironomidae pupae	1	5		Ostracoda	2	10
	Copepoda	1	5		Calanoida	1	5
	Plecoptera adult	1	5		Cyclopoida	1	5
	Subtotal:	601	3,005		Subtotal:	2,573	12,865
	Subtotal all depths for this sampling area and date period:	7,550	12,583	13 Sep 95 9 m	No. samples processed:	4	
					Oligochaeta	3,337	16,685

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
	Chironomidae larvae	132	660		Subtotal all depths for this sampling area and date period:	14,236	23,727
	Cyclopoida	6	30		Total all dates and depths for Silcott Island:	135,211	12,557
	Chironomidae pupae	4	20				
	Bivalvia	3	15				
	Ostracoda	3	15	Centennial Island			
	Nematoda	2	10	30 Mar 94			
	Chironomidae adult	1	5	3 m	No. samples processed:	4	
	Nemertea	1	5		Oligochaeta	1,472	7,360
	Polychaeta	1	5		Chironomidae larvae	299	1,495
	Subtotal:	3,490	17,450		Bivalvia	46	230
13 Sep 95					Eggs (unidentified)	21	105
18 m	No. samples processed:	4			Prostigmata (water mites)	8	40
	Oligochaeta	909	4,545		Nematoda	7	35
	Chironomidae larvae	52	260		Ostracoda	3	15
	Bivalvia	3	15		Cladocera	2	10
	Chironomidae pupae	1	5		Chironomidae pupae	1	5
	Ephemeroptera nymph	1	5		<i>Corophium spinicorne</i>	1	5
	Subtotal:	966	4,830		Ephemeroptera nymph	1	5
	Subtotal all depths for this sampling area and date period:	7,029	11,715		Subtotal:	1,861	9,305
11 Oct 95				30 Mar 94			
3 m	No. samples processed:	4		9 m	No. samples processed:	4	
	Oligochaeta	9,433	47,165		Oligochaeta	1,425	7,125
	Chironomidae larvae	654	3,270		Chironomidae larvae	223	1,115
	Cyclopoida	11	55		Eggs (unidentified)	95	475
	Nematoda	9	45		Nematoda	11	55
	Chironomidae pupae	7	35		Turbellaria	11	55
	Ostracoda	7	35		Cladocera	8	40
	Copepoda	6	30		Prostigmata (water mites)	4	20
	Harpacticoida	4	20		Harpacticoida	3	15
	Bivalvia	3	15		Ostracoda	3	15
	Ephemeroptera nymph	3	15		Copepoda	2	10
	Subtotal:	10,137	50,685		<i>Corophium salmonis</i>	2	10
11 Oct 95					Bivalvia	1	5
9 m	No. samples processed:	4			Cyclopoida	1	5
	Oligochaeta	1,890	9,450		Subtotal:	1,789	8,945
	Chironomidae larvae	119	595	30 Mar 94			
	Copepoda	5	25	18 m	No. samples processed:	4	
	Bivalvia	3	15		Oligochaeta	2,046	10,230
	Chironomidae pupae	1	5		Chironomidae larvae	270	1,350
	Nemertea	1	5		Eggs (unidentified)	80	400
	Subtotal:	2,019	10,095		Cladocera	54	270
11 Oct 95					Copepoda	39	195
18 m	No. samples processed:	4			Nematoda	4	20
	Oligochaeta	1,998	9,990		Nemertea	4	20
	Chironomidae larvae	68	340		Bivalvia	3	15
	Bivalvia	13	65		Chironomidae pupae	3	15
	Copepoda	1	5		Harpacticoida	3	15
	Subtotal:	2,080	10,400		Calanoida	2	10
					Ostracoda	2	10

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
	Plecoptera	2	10	1 Jun 94			
	Araneae (spiders)	1	5	3 m	No. samples processed:	4	
	Cyclopoida	1	5		Chironomidae larvae	163	815
	Plecoptera nymph	1	5		Oligochaeta	147	735
	Trichoptera larvae	1	5		Bivalvia	67	335
	Subtotal:	2,516	12,580		Prostigmata (water mites)	16	80
	Subtotal all depths for this sampling area and date period:	6,166	10,277		Nematoda	12	60
4 May 94					Chironomidae pupae	7	35
3 m	No. samples processed:	4			Chironominae pupae	3	15
	Oligochaeta	471	2,355		Eggs (unidentified)	3	15
	Chironomidae larvae	72	360		<i>Ramelligammarus oregonensis</i>	3	15
	Ostracoda	29	145		Cyclopoida	2	10
	Bivalvia	12	60		Ostracoda	2	10
	Nematoda	12	60		Trichoptera larvae	2	10
	Eggs (unidentified)	3	15		Collembola	1	5
	Prostigmata (water mites)	2	10		Diptera larvae	1	5
	Trichoptera larvae	2	10		Nemertea	1	5
	Copepoda	1	5		Subtotal:	430	2,150
	Ephemeroptera nymph	1	5	1 Jun 94			
	Psychomyiidae larvae	1	5	9 m	No. samples processed:	4	
	Subtotal:	606	3,030		Oligochaeta	1,025	5,125
4 May 94					Chironomidae larvae	321	1,605
9 m	No. samples processed:	4			Nematoda	45	225
	Oligochaeta	1,010	5,050		Harpacticoida	41	205
	Chironomidae larvae	228	1,140		Ostracoda	8	40
	Eggs (unidentified)	34	170		Turbellaria	5	25
	Nematoda	9	45		Prostigmata (water mites)	2	10
	Chironomidae pupae	5	25		Bivalvia	1	5
	Ostracoda	5	25		Calanoida	1	5
	Cyclopoida	3	15		Subtotal:	1,449	7,245
	Bivalvia	2	10	1 Jun 94			
	<i>Corophium salmonis</i>	1	5	18 m	No. samples processed:	4	
	Ephemeroptera nymph	1	5		Oligochaeta	668	3,340
	Subtotal:	1,298	6,490		Chironomidae larvae	179	895
4 May 94					Nematoda	18	90
18 m	No. samples processed:	4			Nemertea	17	85
	Oligochaeta	757	3,785		Cyclopoida	4	20
	Chironomidae larvae	80	400		Harpacticoida	4	20
	Nemertea	32	160		Cladocera	3	15
	Cladocera	14	70		Turbellaria	3	15
	Nematoda	5	25		Chironomidae pupae	1	5
	Cyclopoida	3	15		Ostracoda	1	5
	Ostracoda	1	5		Prostigmata (water mites)	1	5
	Prostigmata (water mites)	1	5		Subtotal:	899	4,495
	Subtotal:	893	4,465		Subtotal all depths for this sampling area and date period:	2,778	4,630
	Subtotal all depths for this sampling area and date period:	2,797	4,662	29 Jun 94			
				3 m	No. samples processed:	4	
					Oligochaeta	387	1,935

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
	Chironomidae larvae	227	1,135		Calanoida	4	20
	Bivalvia	14	70		Chironomidae pupae	4	20
	Chironomidae pupae	7	35		Ephemeroptera nymph	1	5
	Nematoda	6	30		Ostracoda	1	5
	Eggs (unidentified)	5	25		Subtotal:	419	2,095
	Cladocera	2	10	27 Jul 94			
	Harpacticoida	2	10	9 m	No. samples processed:	4	
	Prostigmata (water mites)	2	10		Oligochaeta	301	1,505
	Ephemeroptera nymph	1	5		Chironomidae larvae	53	265
	Nemertea	1	5		Cyclopoida	7	35
	Orthoptera adult	1	5		Nematoda	4	20
	Trichoptera larvae	1	5		Chironomidae adult	2	10
	Subtotal:	656	3,280		Chironomidae pupae	1	5
29 Jun 94					Subtotal:	368	1,840
9 m	No. samples processed:	4		27 Jul 94			
	Oligochaeta	282	1,410	18 m	No. samples processed:	4	
	Chironomidae larvae	70	350		Oligochaeta	515	2,575
	Nematoda	31	155		Chironomidae larvae	99	495
	Harpacticoida	17	85		Cyclopoida	18	90
	Chironomidae pupae	1	5		Calanoida	2	10
	Cladocera	1	5		Chironomidae adult	2	10
	Culicidae adult	1	5		Chironomidae pupae	2	10
	Prostigmata (water mites)	1	5		Nematoda	1	5
	Subtotal:	404	2,020		Trichoptera larvae	1	5
29 Jun 94					Subtotal:	640	3,200
18 m	No. samples processed:	4			Subtotal all depths for this sampling area and date period:	1,427	2,378
	Oligochaeta	296	1,480	24 Aug 94			
	Chironomidae larvae	98	490	3 m	No. samples processed:	4	
	Eggs (unidentified)	43	215		Oligochaeta	844	4,220
	Harpacticoida	21	105		Chironomidae larvae	545	2,725
	Cyclopoida	4	20		Cladocera	464	2,320
	Cladocera	3	15		Bivalvia	64	320
	Ostracoda	3	15		Leptodora kindtii	50	250
	Nematoda	2	10		Eggs (unidentified)	48	240
	Bivalvia	1	5		Copepoda	37	185
	Diptera adult	1	5		Cyclopoida	36	180
	Prostigmata (water mites)	1	5		Chironomidae pupae	18	90
	Trichoptera larvae	1	5		Ostracoda	17	85
	Subtotal:	474	2,370		Leptodoridae	16	80
	Subtotal all depths for this sampling area and date period:	1,534	2,557		Nemertea	16	80
27 Jul 94					Nematoda	11	55
3 m	No. samples processed:	4			Turbellaria	4	20
	Oligochaeta	207	1,035		Prostigmata (water mites)	3	15
	Chironomidae larvae	118	590		Chironominae pupae	1	5
	Cyclopoida	56	280		Ephemeroptera nymph	1	5
	Bivalvia	13	65		Trichoptera larvae	1	5
	Nematoda	9	45		Subtotal:	2,176	10,880
	Prostigmata (water mites)	6	30				

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
24 Aug 94 9 m	No. samples processed:	4			Calanoida	31	155
	Cladocera	1,865	9,325		Bivalvia	19	95
	Oligochaeta	721	3,605		Nematoda	18	90
	<i>Leptodora kindtii</i>	189	945		Mysidacea	5	25
	Chironomidae larvae	185	925		Chironomidae pupae	3	15
	Cyclopoida	125	625		Harpacticoida	2	10
	Copepoda	20	100		Hirudinea	2	10
	Harpacticoida	16	80		Prostigmata (water mites)	2	10
	Ostracoda	10	50		Ceratopogonidae larvae	1	5
	Chironomidae pupae	4	20		Polychaeta	1	5
	Nematoda	3	15		Unidentified	1	5
	Calanoida	2	10		Subtotal:	1,827	9,135
	Orthocladinae pupae	1	5	21 Sep 94 18 m	No. samples processed:	4	
	Subtotal:	3,141	15,705		Cladocera	1,911	9,555
24 Aug 94 18 m	No. samples processed:	4			Oligochaeta	1,526	7,630
	Oligochaeta	1,046	5,230		Cyclopoida	286	1,430
	Chironomidae larvae	162	810		<i>Leptodora kindtii</i>	141	705
	Copepoda	74	370		Chironomidae larvae	127	635
	Cyclopoida	14	70		Calanoida	48	240
	Chironomidae pupae	6	30		Nematoda	11	55
	Ostracoda	5	25		Ostracoda	3	15
	Nemertea	3	15		Chironominae pupae	2	10
	Ephemeroptera nymph	1	5		Subtotal:	4,055	20,275
	Nematoda	1	5		Subtotal all depths for this sampling area and date period:	6,805	11,342
	Subtotal:	1,312	6,560	26 Oct 94 3 m	No. samples processed:	4	
	Subtotal all depths for this sampling area and date period:	6,629	11,048		Oligochaeta	138	690
21 Sep 94 3 m	No. samples processed:	4			Chironomidae larvae	92	460
	Chironomidae larvae	388	1,940		Bivalvia	83	415
	Cyclopoida	188	940		Cyclopoida	19	95
	Oligochaeta	185	925		Prostigmata (water mites)	10	50
	Bivalvia	105	525		Trichoptera larvae	6	30
	Nematoda	19	95		Ephemeroptera nymph	3	15
	Prostigmata (water mites)	15	75		Chironomidae adult	1	5
	Calanoida	11	55		Subtotal:	352	1,760
	Ostracoda	6	30	26 Oct 94 9 m	No. samples processed:	4	
	Ephemeroptera nymph	2	10		Oligochaeta	1,032	5,160
	Trichoptera larvae	2	10		Chironomidae larvae	154	770
	Chironomidae pupae	1	5		Cyclopoida	20	100
	Collembola	1	5		Nematoda	11	55
	Subtotal:	923	4,615		Bivalvia	4	20
21 Sep 94 9 m	No. samples processed:	4			Chironomidae pupae	1	5
	Oligochaeta	1,098	5,490		<i>Corophium spinicorne</i>	1	5
	Chironomidae larvae	339	1,695		Ostracoda	1	5
	Cyclopoida	305	1,525		Prostigmata (water mites)	1	5
					Subtotal:	1,225	6,125

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
26 Oct 94							
18 m	No. samples processed:	4			Subtotal all depths for this sampling area and date period:	4,974	8,290
	Oligochaeta	1,038	5,190	18 Jan 95			
	Chironomidae larvae	123	615	3 m	No. samples processed:	4	
	Homoptera adult	4	20		Oligochaeta	971	4,855
	Cyclopoida	2	10		Chironomidae larvae	168	840
	Bivalvia	1	5		Bivalvia	21	105
	Harpacticoida	1	5		Prostigmata (water mites)	8	40
	Nematoda	1	5		Ostracoda	6	30
	Subtotal:	1,170	5,850		Nematoda	5	25
	Subtotal all depths for this sampling area and date period:	2,747	4,578		Cyclopoida	4	20
30 Nov 94					<i>Corophium</i> spp.	1	5
3 m	No. samples processed:	4			Ephemeroptera nymph	1	5
	Oligochaeta	828	4,140		Psocoptera	1	5
	Chironomidae larvae	265	1,325		Subtotal:	1,186	5,930
	Bivalvia	178	890	18 Jan 95			
	Copepoda	59	295	9 m	No. samples processed:	4	
	Prostigmata (water mites)	54	270		Oligochaeta	871	4,355
	Cyclopoida	10	50		Chironomidae larvae	144	720
	Psychomyiidae larvae	8	40		Bivalvia	8	40
	<i>Caenis</i> spp.	6	30		Prostigmata (water mites)	5	25
	Ostracoda	5	25		Nematoda	4	20
	Ephemeroptera nymph	3	15		<i>Corophium</i> spp.	2	10
	Nematoda	3	15		Ostracoda	2	10
	Gastropoda	1	5		Calanoida	1	5
	Subtotal:	1,420	7,100		Cyclopoida	1	5
					Subtotal:	1,038	5,190
30 Nov 94				18 Jan 95			
9 m	No. samples processed:	4		18 m	No. samples processed:	4	
	Oligochaeta	1,308	6,540		Oligochaeta	2,204	11,020
	Chironomidae larvae	242	1,210		Chironomidae larvae	310	1,550
	Bivalvia	49	245		Cyclopoida	2	10
	Copepoda	20	100		Nematoda	2	10
	Ostracoda	6	30		Ostracoda	2	10
	Ephemeroptera nymph	2	10		Subtotal:	2,520	12,600
	Nematoda	2	10		Subtotal all depths for this sampling area and date period:	4,744	7,907
	Prostigmata (water mites)	2	10	22 Mar 95			
	Cladocera	1	5	3 m	No. samples processed:	4	
	Hirudinea	1	5		Oligochaeta	429	2,145
	Subtotal:	1,633	8,165		Chironomidae larvae	132	660
30 Nov 94					Bivalvia	70	350
18 m	No. samples processed:	4			Prostigmata (water mites)	19	95
	Oligochaeta	1,549	7,745		Ephemeroptera nymph	15	75
	Chironomidae larvae	353	1,765		Ostracoda	11	55
	Copepoda	7	35		Cyclopoida	5	25
	Ostracoda	7	35		Nemertea	2	10
	Bivalvia	3	15		Nematoda	1	5
	Ephemeroptera nymph	2	10		Trichoptera larvae	1	5
	Subtotal:	1,921	9,605				

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
	Turbellaria	1	5		Nemertea	11	55
	Subtotal:	686	3,430		Ostracoda	10	50
22 Mar 95					Prostigmata (water mites)	5	25
9 m	No. samples processed:	4			Turbellaria	5	25
	Oligochaeta	825	4,125		Bivalvia	2	10
	Chironomidae larvae	123	615		Ephemeroptera nymph	1	5
	Nematoda	31	155		Trichoptera larvae	1	5
	Nemertea	7	35		Subtotal:	2,478	12,390
	Prostigmata (water mites)	3	15	26 Apr 95			
	Cyclopoida	2	10	18 m	No. samples processed:	4	
	Ostracoda	2	10		Oligochaeta	2,692	13,460
	Bivalvia	1	5		Chironomidae larvae	243	1,215
	<i>Corophium</i> spp.	1	5		Ostracoda	34	170
	Subtotal:	995	4,975		Cyclopoida	18	90
22 Mar 95					Nematoda	9	45
18 m	No. samples processed:	4			Prostigmata (water mites)	3	15
	Oligochaeta	1,648	8,240		Turbellaria	3	15
	Chironomidae larvae	218	1,090		Harpacticoida	2	10
	Nematoda	7	35		Ephemeroptera nymph	1	5
	Turbellaria	2	10		Nemertea	1	5
	Cyclopoida	1	5		Subtotal:	3,006	15,030
	Subtotal:	1,876	9,380		Subtotal all depths for this sampling area and date period:	6,773	11,288
	Subtotal all depths for this sampling area and date period:	3,557	5,928	24 May 95			
26 Apr 95				3 m	No. samples processed:	4	
3 m	No. samples processed:	4			Oligochaeta	1,206	6,030
	Oligochaeta	824	4,120		Chironomidae larvae	122	610
	Chironomidae larvae	225	1,125		Bivalvia	118	590
	Bivalvia	177	885		Harpacticoida	21	105
	Prostigmata (water mites)	21	105		Nematoda	20	100
	Ostracoda	12	60		Nemertea	12	60
	Ephemeroptera nymph	10	50		Chironomidae pupae	6	30
	Nemertea	8	40		Cyclopoida	5	25
	Nematoda	3	15		Ostracoda	5	25
	Copepoda	2	10		Prostigmata (water mites)	5	25
	Trichoptera larvae	2	10		Ephemeroptera nymph	4	20
	Araneae (spiders)	1	5		<i>Corophium</i> spp.	2	10
	Chironomidae pupae	1	5		Trichoptera larvae	2	10
	Collembola	1	5		Subtotal:	1,528	7,640
	Cyclopoida	1	5	24 May 95			
	Polychaeta	1	5	9 m	No. samples processed:	4	
	Subtotal:	1,289	6,445		Oligochaeta	2,290	11,450
26 Apr 95					Chironomidae larvae	293	1,465
9 m	No. samples processed:	4			Nematoda	95	475
	Oligochaeta	2,068	10,340		Harpacticoida	59	295
	Chironomidae larvae	244	1,220		Copepoda	19	95
	Harpacticoida	71	355		Nemertea	14	70
	Nematoda	42	210		Ostracoda	5	25
	Cyclopoida	18	90		Prostigmata (water mites)	4	20

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
	Cyclopoida	2	10		Chironomidae pupae	1	5
	Bivalvia	1	5		<i>Corophium salmonis</i>	1	5
	Ephemeroptera nymph	1	5		Hemiptera adult	1	5
	Hemiptera	1	5		Subtotal:	1,892	9,460
	Turbellaria	1	5	21 Jun 95			
	Subtotal:	2,785	13,925	18 m	No. samples processed:	4	
24 May 95					Oligochaeta	1,476	7,380
18 m	No. samples processed:	4			Chironomidae larvae	211	1,055
	Oligochaeta	2,923	14,615		Nematoda	13	65
	Chironomidae larvae	388	1,940		Harpacticoida	4	20
	Nematoda	173	865		Nemertea	3	15
	Harpacticoida	35	175		Chironomidae pupae	1	5
	Cyclopoida	22	110		Cyclopoida	1	5
	Ostracoda	8	40		Ostracoda	1	5
	Nemertea	7	35		Prostigmata (water mites)	1	5
	Collembola	6	30		Trichoptera larvae	1	5
	Ephemeroptera nymph	4	20		Subtotal:	1,712	8,560
	Bivalvia	3	15		Subtotal all depths for this sampling area and date period:	4,377	7,295
	Ceratopogonidae larvae	1	5	19 Jul 95			
	Diptera	1	5	3 m	No. samples processed:	4	
	Prostigmata (water mites)	1	5		Oligochaeta	804	4,020
	Subtotal:	3,572	17,860		Chironomidae larvae	254	1,270
	Subtotal all depths for this sampling area and date period:	7,885	13,142		Bivalvia	33	165
21 Jun 95					Nematoda	14	70
3 m	No. samples processed:	4			Chironomidae pupae	10	50
	Oligochaeta	645	3,225		Prostigmata (water mites)	8	40
	Chironomidae larvae	99	495		Ostracoda	5	25
	Bivalvia	19	95		Trichoptera larvae	3	15
	Prostigmata (water mites)	3	15		Cyclopoida	2	10
	Chironomidae pupae	2	10		Harpacticoida	2	10
	Nematoda	2	10		Copepoda	1	5
	Ephemeroptera nymph	1	5		Diptera	1	5
	Harpacticoida	1	5		Nemertea	1	5
	<i>Porcellio</i> spp.	1	5		Subtotal:	1,138	5,690
	Subtotal:	773	3,865	19 Jul 95			
21 Jun 95				9 m	No. samples processed:	4	
9 m	No. samples processed:	4			Oligochaeta	1,212	6,060
	Oligochaeta	1,525	7,625		Chironomidae larvae	367	1,835
	Chironomidae larvae	228	1,140		Nematoda	26	130
	Nematoda	72	360		Harpacticoida	9	45
	Harpacticoida	39	195		Ostracoda	5	25
	Nemertea	8	40		Bivalvia	2	10
	Cyclopoida	7	35		Cyclopoida	2	10
	Turbellaria	3	15		Copepoda	1	5
	Collembola	2	10		Diptera	1	5
	Ixodides (ticks)	2	10		Nemertea	1	5
	Prostigmata (water mites)	2	10		Prostigmata (water mites)	1	5
	Bivalvia	1	5		Subtotal:	1,627	8,135

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
19 Jul 95							
18 m	No. samples processed:	4				Subtotal:	828 4,140
	Oligochaeta	1,131	5,655			Subtotal all depths for this sampling area and date period:	4,611 7,685
	Chironomidae larvae	291	1,455	13 Sep 95			
	Nematoda	4	20	3 m	No. samples processed:	4	
	Ostracoda	4	20		Oligochaeta	442	2,210
	Bivalvia	3	15		Bivalvia	72	360
	Copepoda	3	15		Chironomidae larvae	32	160
	Nemertea	1	5		Ostracoda	8	40
	Prostigmata (water mites)	1	5		Prostigmata (water mites)	5	25
	Subtotal:	1,438	7,190		Collembola	2	10
	Subtotal all depths for this sampling area and date period:	4,203	7,005		<i>Corophium spinicorne</i>	2	10
16 Aug 95					Chironomidae pupae	1	5
3 m	No. samples processed:	4			Copepoda	1	5
	Oligochaeta	882	4,410		Diptera adult	1	5
	Chironomidae larvae	179	895		Ephemeroptera nymph	1	5
	Bivalvia	36	180		Nematoda	1	5
	Harpacticoida	10	50		Subtotal:	568	2,840
	Ostracoda	10	50	13 Sep 95			
	Nematoda	5	25	9 m	No. samples processed:	4	
	Prostigmata (water mites)	5	25		Oligochaeta	1,031	5,155
	Chironomidae pupae	3	15		Chironomidae larvae	15	75
	Cyclopoida	3	15		Cyclopoida	3	15
	Polychaeta	1	5		Bivalvia	2	10
	Subtotal:	1,134	5,670		Copepoda	2	10
16 Aug 95					Ostracoda	2	10
9 m	No. samples processed:	4			Harpacticoida	1	5
	Oligochaeta	2,435	12,175		Nematoda	1	5
	Chironomidae larvae	188	940		Subtotal:	1,057	5,285
	Nematoda	6	30	13 Sep 95			
	Ostracoda	5	25	18 m	No. samples processed:	4	
	Harpacticoida	3	15		Oligochaeta	394	1,970
	Nemertea	3	15		Chironomidae larvae	4	20
	Copepoda	2	10		Collembola	2	10
	Cyclopoida	2	10		Nemertea	1	5
	Hirudinea	2	10		Ostracoda	1	5
	Bivalvia	1	5		Subtotal:	402	2,010
	Chironomidae pupae	1	5		Subtotal all depths for this sampling area and date period:	2,027	3,378
	Plecoptera nymph	1	5	11 Oct 95			
	Subtotal:	2,649	13,245	3 m	No. samples processed:	4	
16 Aug 95					Oligochaeta	515	2,575
18 m	No. samples processed:	4			Bivalvia	28	140
	Oligochaeta	788	3,940		Chironomidae larvae	13	65
	Chironomidae larvae	32	160		Ephemeroptera nymph	5	25
	Nematoda	4	20		Ostracoda	2	10
	Nemertea	2	10		Chironomidae pupae	1	5
	<i>Corophium</i> spp.	1	5		Subtotal:	564	2,820
	Ostracoda	1	5	11 Oct 95			
				9 m	No. samples processed:	4	

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
	Oligochaeta	749	3,745		Cyclopoida	3	15
	Chironomidae larvae	15	75		<i>Hexagenia</i> spp.	3	15
	Nematoda	5	25		Nemertea	3	15
	Harpacticoida	2	10		Cladocera	1	5
	Chironomidae pupae	1	5		<i>Corophium spinicorne</i>	1	5
	Copepoda	1	5		Elmidae adult	1	5
	Ephemeroptera nymph	1	5		<i>Ephemera</i> spp.	1	5
	Subtotal:	774	3,870		Trichoptera larvae	1	5
11 Oct 95					Subtotal:	757	3,785
18 m	No. samples processed:	4		30 Mar 94			
	Oligochaeta	937	4,685	18 m	No. samples processed:	4	
	Chironomidae larvae	30	150		Oligochaeta	790	3,950
	Coleoptera adult	1	5		Chironomidae larvae	178	890
	Ephemeroptera nymph	1	5		Bivalvia	43	215
	Subtotal:	969	4,845		Eggs (unidentified)	43	215
	Subtotal all depths for this sampling area and date period:	2,307	3,845		Prostigmata (water mites)	13	65
	Total all dates and depths for Centennial Island:	76,341	7,069		Ostracoda	8	40
					Copepoda	5	25
					Nematoda	4	20
Offield					Cyclopoida	2	10
30 Mar 94					Coleoptera	1	5
3 m	No. samples processed:	4			Nemertea	1	5
	Oligochaeta	297	1,485		Polychaeta	1	5
	Bivalvia	270	1,350		Trichoptera larvae	1	5
	Chironomidae larvae	209	1,045		Subtotal:	1,090	5,450
	Ephemeroptera nymph	49	245		Subtotal all depths for this sampling area and date period:	2,743	4,572
	Prostigmata (water mites)	27	135				
	Eggs (unidentified)	11	55	4 May 94			
	Nematoda	11	55	3 m	No. samples processed:	4	
	Trichoptera larvae	6	30		Oligochaeta	132	660
	Caenidae nymph	5	25		Chironomidae larvae	48	240
	Ceratopogonidae larvae	4	20		Bivalvia	43	215
	<i>Corophium salmonis</i>	2	10		Ostracoda	20	100
	Harpacticoida	2	10		Nemertea	14	70
	Araneae (spiders)	1	5		Ephemeroptera nymph	10	50
	Cyclopoida	1	5		Prostigmata (water mites)	6	30
	<i>Ephemera</i> spp.	1	5		Harpacticoida	5	25
	Subtotal:	896	4,480		Ceratopogonidae larvae	4	20
30 Mar 94					Nematoda	2	10
9 m	No. samples processed:	4			Copepoda	1	5
	Oligochaeta	532	2,660		Cyclopoida	1	5
	Chironomidae larvae	82	410		Subtotal:	286	1,430
	Bivalvia	73	365				
	Prostigmata (water mites)	25	125	4 May 94			
	Nematoda	10	50	9 m	No. samples processed:	4	
	Eggs (unidentified)	9	45		Oligochaeta	158	790
	Ephemeroptera nymph	5	25		Chironomidae larvae	143	715
	Turbellaria	4	20		Bivalvia	12	60
	Copepoda	3	15		Nematoda	12	60
					Chironomidae pupae	1	5

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
	Eggs (unidentified)	1	5		Ostracoda	1	5
	Ephemeroptera nymph	1	5		Subtotal:	162	810
	Subtotal:	328	1,640	1 Jun 94			
4 May 94				18 m	No. samples processed:	4	
18 m	No. samples processed:	4		Oligochaeta		204	1,020
	Oligochaeta	530	2,650	Chironomidae larvae		95	475
	Chironomidae larvae	83	415	Eggs (unidentified)		24	120
	Bivalvia	63	315	Bivalvia		8	40
	Eggs (unidentified)	16	80	Prostigmata (water mites)		4	20
	Cyclopoida	8	40	Nematoda		2	10
	Nemertea	8	40	Ephemeroptera nymph		1	5
	Nematoda	2	10	Nemertea		1	5
	Chironomidae pupae	1	5		Subtotal:	339	1,695
	Ostracoda	1	5		Subtotal all depths for this sampling area and date period:	1,121	1,868
	Prostigmata (water mites)	1	5				
	Subtotal:	713	3,565	29 Jun 94			
	Subtotal all depths for this sampling area and date period:	1,327	2,212	3 m	No. samples processed:	4	
1 Jun 94				Oligochaeta		556	2,780
3 m	No. samples processed:	4		Chironomidae larvae		371	1,855
	Oligochaeta	247	1,235	Bivalvia		156	780
	Chironomidae larvae	187	935	Chironomidae pupae		33	165
	Bivalvia	84	420	Cladocera		22	110
	Nematoda	40	200	Copepoda		19	95
	Ephemeroptera nymph	17	85	Eggs (unidentified)		19	95
	Harpacticoida	13	65	Ephemeroptera nymph		17	85
	Chironomidae pupae	12	60	Prostigmata (water mites)		10	50
	Prostigmata (water mites)	10	50	Cyclopoida		8	40
	Cyclopoida	4	20	<i>Hexagenia</i> spp.		5	25
	Ceratopogonidae larvae	2	10	Isopoda		2	10
	Collembola	1	5	Nematoda		2	10
	<i>Hexagenia</i> spp.	1	5	Orthocladinae pupae		2	10
	Ostracoda	1	5	Trichoptera larvae		2	10
	<i>Ramullogammarus ramellus</i>	1	5	Araneae (spiders)		1	5
	Subtotal:	620	3,100	Ceratopogonidae larvae		1	5
1 Jun 94				Chironominae pupae		1	5
9 m	No. samples processed:	4		Diptera adult		1	5
	Oligochaeta	71	355	Harpacticoida		1	5
	Chironomidae larvae	53	265	Nemertea		1	5
	Bivalvia	14	70	Ostracoda		1	5
	Nematoda	6	30	Polychaeta		1	5
	Cyclopoida	4	20	<i>Ramullogammarus ramellus</i>		1	5
	Copepoda	2	10		Subtotal:	1,233	6,165
	Diptera pupae	2	10	29 Jun 94			
	Eggs (unidentified)	2	10	9 m	No. samples processed:	4	
	Ephemeroptera nymph	2	10	Oligochaeta		270	1,350
	Nemertea	2	10	Chironomidae larvae		104	520
	Prostigmata (water mites)	2	10	Cladocera		26	130
	Chironomidae pupae	1	5	Eggs (unidentified)		22	110
				Bivalvia		17	85

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
	Copepoda	13	65		Chironomidae pupae	4	20
	Nematoda	5	25		Isopoda	2	10
	Diptera adult	4	20		Prostigmata (water mites)	1	5
	Chironomidae pupae	1	5		Trichoptera larvae	1	5
	<i>Corophium spinicorne</i>	1	5		Subtotal:	604	3,020
	Cyclopoida	1	5	27 Jul 94			
	Ephemeroptera nymph	1	5	18 m	No. samples processed:	4	
	Insecta	1	5		Chironomidae larvae	342	1,710
	Prostigmata (water mites)	1	5		Oligochaeta	291	1,455
	Tanypodinae pupae	1	5		Cyclopoida	37	185
	Subtotal:	468	2,340		<i>Corophium spinicorne</i>	36	180
29 Jun 94					Ostracoda	16	80
18 m	No. samples processed:	4			Bivalvia	15	75
	Oligochaeta	320	1,600		Harpacticoida	3	15
	Chironomidae larvae	165	825		Nematoda	3	15
	Cladocera	31	155		Prostigmata (water mites)	2	10
	Bivalvia	12	60		Chironomidae pupae	1	5
	Cyclopoida	8	40		Ephemeroptera nymph	1	5
	Eggs (unidentified)	6	30		Subtotal:	747	3,735
	Nematoda	4	20		Subtotal all depths for this sampling area and date period:	2,526	4,210
	Nemertea	4	20	24 Aug 94			
	Ostracoda	4	20	3 m	No. samples processed:	4	
	Prostigmata (water mites)	2	10		Oligochaeta	499	2,495
	Harpacticoida	1	5		Bivalvia	256	1,280
	Subtotal:	557	2,785		Chironomidae larvae	248	1,240
	Subtotal all depths for this sampling area and date period:	2,258	3,763		Cyclopoida	62	310
27 Jul 94					Nematoda	49	245
3 m	No. samples processed:	4			Prostigmata (water mites)	18	90
	Chironomidae larvae	623	3,115		<i>Hexagenia</i> spp.	16	80
	Oligochaeta	347	1,735		Cladocera	15	75
	Bivalvia	77	385		Ephemeroptera nymph	7	35
	Prostigmata (water mites)	36	180		Ostracoda	6	30
	Nematoda	32	160		<i>Caenis</i> spp.	4	20
	Ephemeroptera nymph	20	100		<i>Leptodora kindtii</i>	4	20
	Cyclopoida	17	85		Chironomidae pupae	3	15
	Chironomidae pupae	15	75		<i>Corophium salmonis</i>	2	10
	Ostracoda	5	25		Harpacticoida	2	10
	Harpacticoida	2	10		Isopoda	2	10
	Trichoptera larvae	1	5		Psychomyiidae larvae	2	10
	Subtotal:	1,175	5,875		Calanoida	1	5
27 Jul 94					Chironominae pupae	1	5
9 m	No. samples processed:	4			Subtotal:	1,197	5,985
	Chironomidae larvae	342	1,710	24 Aug 94			
	Oligochaeta	128	640	9 m	No. samples processed:	4	
	Bivalvia	73	365		Chironomidae larvae	168	840
	Ephemeroptera nymph	19	95		Oligochaeta	162	810
	Cyclopoida	18	90		Bivalvia	28	140
	Nematoda	16	80		Cladocera	21	105

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
	Cyclopoida	16	80		Subtotal:	1,029	5,145
	Ostracoda	11	55	21 Sep 94			
	Nematoda	9	45	9 m	No. samples processed:	4	
	<i>Hexagenia</i> spp.	4	20		Oligochaeta	190	950
	<i>Leptodora kindtii</i>	4	20		Bivalvia	182	910
	Prostigmata (water mites)	3	15		Cladocera	147	735
	Ephemeroptera nymph	2	10		Chironomidae larvae	112	560
	Chironomidae pupae	1	5		Cyclopoida	41	205
	<i>Corophium salmonis</i>	1	5		Nematoda	26	130
	Nemertea	1	5		Calanoida	24	120
	<i>Sialis</i> spp.	1	5		Ostracoda	10	50
	Subtotal:	432	2,160		Ephemeroptera nymph	9	45
24 Aug 94					<i>Hexagenia</i> spp.	4	20
18 m	No. samples processed:	4			Prostigmata (water mites)	4	20
	Chironomidae larvae	458	2,290		<i>Corophium spinicorne</i>	2	10
	Oligochaeta	400	2,000		<i>Leptodora kindtii</i>	2	10
	Cladocera	66	330		<i>Corophium salmonis</i>	1	5
	Ostracoda	46	230		Subtotal:	754	3,770
	Bivalvia	24	120	21 Sep 94			
	Cyclopoida	17	85	18 m	No. samples processed:	4	
	Copepoda	13	65		Oligochaeta	718	3,590
	<i>Leptodora kindtii</i>	5	25		Chironomidae larvae	311	1,555
	Nematoda	5	25		Cyclopoida	148	740
	Chironomidae adult	1	5		Ostracoda	37	185
	Chironomidae pupae	1	5		Bivalvia	32	160
	<i>Corophium spinicorne</i>	1	5		Calanoida	28	140
	Hirudinea	1	5		Nematoda	10	50
	Nemertea	1	5		Prostigmata (water mites)	5	25
	Polychaeta	1	5		Chironomidae pupae	3	15
	Subtotal:	1,040	5,200		<i>Corophium spinicorne</i>	3	15
	Subtotal all depths for this sampling area and date period:	2,669	4,448		Ephemeroptera nymph	1	5
21 Sep 94					Hirudinea	1	5
3 m	No. samples processed:	4			Subtotal:	1,297	6,485
	Bivalvia	405	2,025		Subtotal all depths for this sampling area and date period:	3,080	5,133
	Chironomidae larvae	292	1,460	26 Oct 94			
	Ephemeroptera nymph	125	625	3 m	No. samples processed:	4	
	Oligochaeta	108	540		Oligochaeta	177	885
	Cyclopoida	31	155		Chironomidae larvae	44	220
	<i>Caenis</i> spp.	19	95		Bivalvia	38	190
	Prostigmata (water mites)	17	85		Ephemeroptera nymph	15	75
	Nematoda	12	60		Cyclopoida	6	30
	Calanoida	6	30		Nematoda	2	10
	Chironomidae pupae	5	25		Prostigmata (water mites)	2	10
	Copepoda	4	20		Gastropoda	1	5
	Ostracoda	2	10		Ostracoda	1	5
	Harpacticoida	1	5		Subtotal:	286	1,430
	Leptophlebiidae nymph	1	5	26 Oct 94			
	Trichoptera larvae	1	5	9 m	No. samples processed:	4	

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
	Oligochaeta	45	225		Ephemeroptera nymph	15	75
	Bivalvia	29	145		Nematoda	5	25
	Chironomidae larvae	17	85		Gammaridae	2	10
	Cyclopoida	5	25		Polychaeta	2	10
	<i>Corophium spinicorne</i>	3	15		Cladocera	1	5
	Ephemeroptera nymph	3	15		Nemertea	1	5
	Nematoda	3	15		Subtotal:	541	2,705
	Prostigmata (water mites)	1	5	30 Nov 94			
	Subtotal:	106	530	18 m	No. samples processed:	4	
26 Oct 94					Oligochaeta	161	805
18 m	No. samples processed:	4			Bivalvia	114	570
	Oligochaeta	190	950		Chironomidae larvae	82	410
	Bivalvia	146	730		Copepoda	62	310
	Chironomidae larvae	57	285		Ostracoda	14	70
	Cyclopoida	34	170		<i>Corophium</i> spp.	7	35
	<i>Corophium</i> spp.	9	45		Ephemeroptera nymph	4	20
	Calanoida	2	10		Polychaeta	4	20
	<i>Corophium spinicorne</i>	2	10		Prostigmata (water mites)	3	15
	Ostracoda	2	10		<i>Corophium salmonis</i>	1	5
	Ephemeroptera nymph	1	5		<i>Hyalella azteca</i>	1	5
	Nematoda	1	5		Nematoda	1	5
	Subtotal:	444	2,220		Subtotal:	454	2,270
	Subtotal all depths for this sampling area and date period:	836	1,393		Subtotal all depths for this sampling area and date period:	1,589	2,648
30 Nov 94				18 Jan 95			
3 m	No. samples processed:	4		3 m	No. samples processed:	4	
	Oligochaeta	228	1,140		Oligochaeta	708	3,540
	Bivalvia	143	715		Bivalvia	310	1,550
	Chironomidae larvae	75	375		Chironomidae larvae	231	1,155
	Ephemeroptera nymph	42	210		Cyclopoida	56	280
	Cyclopoida	37	185		Ephemeroptera nymph	45	225
	Prostigmata (water mites)	32	160		Prostigmata (water mites)	33	165
	Nemertea	14	70		Nemertea	14	70
	Nematoda	8	40		Nematoda	13	65
	Ostracoda	6	30		Ostracoda	6	30
	Unidentified	5	25		Harpacticoida	2	10
	<i>Caenis</i> spp.	2	10		Subtotal:	1,418	7,090
	Copepoda	1	5	18 Jan 95			
	Trichoptera larvae	1	5	9 m	No. samples processed:	4	
	Subtotal:	594	2,970		Oligochaeta	676	3,380
30 Nov 94					Bivalvia	307	1,535
9 m	No. samples processed:	4			Cyclopoida	76	380
	Oligochaeta	145	725		Chironomidae larvae	38	190
	Bivalvia	132	660		Prostigmata (water mites)	26	130
	Ostracoda	73	365		<i>Corophium</i> spp.	23	115
	Chironomidae larvae	56	280		Ostracoda	23	115
	Copepoda	55	275		Ephemeroptera nymph	16	80
	<i>Corophium spinicorne</i>	37	185		Nematoda	16	80
	Prostigmata (water mites)	17	85		Nemertea	2	10

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
	Polychaeta	2	10		<i>Hexagenia</i> spp.	1	5
	Chironomidae pupae	1	5		Ostracoda	1	5
	<i>Corophium salmonis</i>	1	5		Trichoptera larvae	1	5
	Turbellaria	1	5		Subtotal:	666	3,330
	Subtotal:	1,208	6,040	22 Mar 95			
18 Jan 95				18 m	No. samples processed:	4	
18 m	No. samples processed:	4			Oligochaeta	244	1,220
	Oligochaeta	721	3,605		Bivalvia	111	555
	Chironomidae larvae	172	860		Chironomidae larvae	25	125
	Bivalvia	125	625		Copepoda	14	70
	Cyclopoida	107	535		Cyclopoida	13	65
	Copepoda	35	175		<i>Corophium</i> spp.	9	45
	Prostigmata (water mites)	24	120		Ostracoda	7	35
	<i>Corophium</i> spp.	17	85		Prostigmata (water mites)	4	20
	Ostracoda	10	50		Chironomidae pupae	1	5
	<i>Corophium salmonis</i>	9	45		Polychaeta	1	5
	Gammaridae	2	10		Subtotal:	429	2,145
	Nematoda	2	10		Subtotal all depths for this sampling area and date period:	1,858	3,097
	Hirudinea	1	5	26 Apr 95			
	Nemertea	1	5	3 m	No. samples processed:	4	
	Subtotal:	1,226	6,130		Oligochaeta	811	4,055
	Subtotal all depths for this sampling area and date period:	3,852	6,420		Bivalvia	201	1,005
22 Mar 95					Chironomidae larvae	126	630
3 m	No. samples processed:	4			<i>Corophium salmonis</i>	123	615
	Oligochaeta	325	1,625		Ephemeroptera nymph	40	200
	Chironomidae larvae	259	1,295		Prostigmata (water mites)	33	165
	Bivalvia	106	530		Nematoda	28	140
	Prostigmata (water mites)	21	105		Copepoda	8	40
	Nematoda	15	75		Ostracoda	7	35
	Ephemeroptera nymph	14	70		Cyclopoida	3	15
	Cyclopoida	11	55		Chironomidae pupae	2	10
	Trichoptera larvae	6	30		Turbellaria	2	10
	Ostracoda	2	10		Ceratopogonidae larvae	1	5
	Coleoptera larvae	1	5		<i>Corophium</i> spp.	1	5
	<i>Corophium</i> spp.	1	5		Subtotal:	1,386	6,930
	Nemertea	1	5	26 Apr 95			
	Polychaeta	1	5	9 m	No. samples processed:	4	
	Subtotal:	763	3,815		Oligochaeta	524	2,620
22 Mar 95					Bivalvia	198	990
9 m	No. samples processed:	4			Chironomidae larvae	102	510
	Oligochaeta	365	1,825		<i>Corophium</i> spp.	86	430
	Bivalvia	121	605		Nematoda	32	160
	Nematoda	71	355		Ephemeroptera nymph	10	50
	Cyclopoida	47	235		Prostigmata (water mites)	7	35
	Chironomidae larvae	36	180		Copepoda	5	25
	Prostigmata (water mites)	12	60		Cyclopoida	5	25
	Ephemeroptera nymph	10	50		Ostracoda	5	25
	<i>Corophium salmonis</i>	1	5		Polychaeta	3	15

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
	Chironomidae pupae	1	5		Subtotal:	756	3,780
	Nemertea	1	5	24 May 95			
	Trichoptera larvae	1	5	18 m	No. samples processed:	4	
	Subtotal:	980	4,900		Oligochaeta	549	2,745
26 Apr 95					Chironomidae larvae	232	1,160
18 m	No. samples processed:	4			Bivalvia	86	430
	Oligochaeta	1,140	5,700		Ostracoda	13	65
	Chironomidae larvae	191	955		Chironomidae pupae	2	10
	Bivalvia	81	405		Copepoda	1	5
	Ostracoda	13	65		Nematoda	1	5
	Copepoda	10	50		Prostigmata (water mites)	1	5
	Cyclopoida	5	25		Turbellaria	1	5
	Polychaeta	4	20		Subtotal:	886	4,430
	Prostigmata (water mites)	4	20		Subtotal all depths for this sampling area and date period:	2,360	3,933
	<i>Corophium salmons</i>	2	10	21 Jun 95			
	Nematoda	2	10	3 m	No. samples processed:	4	
	<i>Corophium</i> spp.	1	5		Oligochaeta	187	935
	Ephemeroptera nymph	1	5		Chironomidae larvae	93	465
	Subtotal:	1,454	7,270		Bivalvia	38	190
	Subtotal all depths for this sampling area and date period:	3,820	6,367		Nematoda	5	25
24 May 95					Cyclopoida	4	20
3 m	No. samples processed:	4			Ostracoda	2	10
	Oligochaeta	515	2,575		Prostigmata (water mites)	2	10
	Bivalvia	68	340		Chironomidae pupae	1	5
	Chironomidae larvae	53	265		Ephemeroptera nymph	1	5
	Nematoda	45	225		<i>Hexagenia</i> spp.	1	5
	Prostigmata (water mites)	11	55		Subtotal:	334	1,670
	Cyclopoida	10	50	21 Jun 95			
	Ephemeroptera nymph	5	25	9 m	No. samples processed:	4	
	Chironomidae pupae	4	20		Oligochaeta	57	285
	Ostracoda	3	15		Chironomidae larvae	18	90
	Trichoptera larvae	3	15		Bivalvia	4	20
	Nemertea	1	5		Ostracoda	4	20
	Subtotal:	718	3,590		Nematoda	3	15
24 May 95					Cyclopoida	1	5
9 m	No. samples processed:	4			Ephemeroptera nymph	1	5
	Oligochaeta	450	2,250		Subtotal:	88	440
	Chironomidae larvae	126	630	21 Jun 95			
	Bivalvia	78	390	18 m	No. samples processed:	4	
	Nematoda	65	325		Oligochaeta	1,159	5,795
	Copepoda	10	50		Chironomidae larvae	177	885
	Ephemeroptera nymph	10	50		Bivalvia	18	90
	Ostracoda	5	25		Nematoda	6	30
	Prostigmata (water mites)	4	20		Ostracoda	3	15
	<i>Corophium salmons</i>	3	15		Cyclopoida	1	5
	Chironomidae pupae	2	10		Prostigmata (water mites)	1	5
	<i>Corophium</i> spp.	2	10		Subtotal:	1,365	6,825
	Collembola	1	5				

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
Subtotal all depths for this sampling area and date period:				Thysanoptera adult			
		1,787	2,978			2	10
19 Jul 95					Cyclopoida	1	5
3 m	No. samples processed:	4			Harpacticoida	1	5
	Oligochaeta	492	2,460		Ostracoda	1	5
	Chironomidae larvae	327	1,635		Trichoptera adult	1	5
	Bivalvia	50	250		Subtotal:	820	4,100
	Nematoda	25	125		Subtotal all depths for this sampling area and date period:	2,494	4,157
	Copepoda	13	65	16 Aug 95			
	Ephemeroptera nymph	12	60	3 m	No. samples processed:	4	
	Ostracoda	6	30		Oligochaeta	94	470
	Prostigmata (water mites)	6	30		Chironomidae larvae	78	390
	Cyclopoida	3	15		Copepoda	14	70
	Chironomidae adult	1	5		Bivalvia	7	35
	Chironomidae pupae	1	5		Ostracoda	3	15
	Hirudinea	1	5		<i>Corophium</i> spp.	2	10
	Homoptera adult	1	5		Cyclopoida	2	10
	Polychaeta	1	5		Nematoda	2	10
	Thysanoptera adult	1	5		Subtotal:	202	1,010
	Subtotal:	940	4,700	16 Aug 95			
19 Jul 95				9 m	No. samples processed:	4	
9 m	No. samples processed:	4			Oligochaeta	817	4,085
	Oligochaeta	419	2,095		Chironomidae larvae	242	1,210
	Chironomidae larvae	199	995		Cyclopoida	17	85
	Bivalvia	46	230		Bivalvia	9	45
	Nematoda	22	110		Ephemeroptera nymph	3	15
	Cyclopoida	13	65		Nematoda	3	15
	Copepoda	8	40		Ostracoda	3	15
	Homoptera adult	8	40		<i>Corophium</i> spp.	2	10
	Ephemeroptera nymph	7	35		Chironomidae pupae	1	5
	Prostigmata (water mites)	4	20		Copepoda	1	5
	Trichoptera larvae	3	15		Polychaeta	1	5
	<i>Corophium salmonis</i>	2	10		Prostigmata (water mites)	1	5
	Ostracoda	2	10		Subtotal:	1,100	5,500
	Polychaeta	1	5	16 Aug 95			
	Subtotal:	734	3,670	18 m	No. samples processed:	4	
19 Jul 95					Oligochaeta	1,424	7,120
18 m	No. samples processed:	4			Chironomidae larvae	157	785
	Oligochaeta	685	3,425		Copepoda	45	225
	Chironomidae larvae	74	370		Turbellaria	3	15
	Homoptera adult	24	120		Nematoda	2	10
	<i>Corophium salmonis</i>	8	40		Chironomidae pupae	1	5
	Bivalvia	5	25		Subtotal:	1,632	8,160
	Chironomidae adult	4	20		Subtotal all depths for this sampling area and date period:	2,934	4,890
	Hemiptera	4	20	13 Sep 95			
	Nematoda	4	20	3 m	No. samples processed:	4	
	Chironomidae pupae	2	10		Oligochaeta	157	785
	Coleoptera adult	2	10		Chironomidae larvae	49	245
	Diptera adult	2	10				

Appendix Table C1. Continued.

Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)	Sampling area Date Depth (m)	Taxon/category	Total number	Density (no./m ²)
	Bivalvia	24	120		Nematoda	21	105
	Ephemeroptera nymph	5	25		Prostigmata (water mites)	3	15
	Chironomidae pupae	4	20		Ostracoda	2	10
	Prostigmata (water mites)	4	20		Polychaeta	2	10
	<i>Corophium salmonis</i>	2	10		Copepoda	1	5
	Ostracoda	2	10		Subtotal:	1,079	5,395
	Nematoda	1	5	11 Oct 95			
	Polychaeta	1	5	18 m	No. samples processed:	4	
	Trichoptera larvae	1	5		Oligochaeta	1,048	5,240
	Subtotal:	250	1,250		Chironomidae larvae	100	500
13 Sep 95					Bivalvia	9	45
9 m	No. samples processed:	4			Homoptera adult	3	15
	Oligochaeta	180	900		Nematoda	3	15
	Chironomidae larvae	30	150		Ostracoda	3	15
	Polychaeta	8	40		Diptera adult	2	10
	Nematoda	6	30		Polychaeta	2	10
	Bivalvia	4	20		Prostigmata (water mites)	2	10
	Chironomidae adult	1	5		Chironomidae adult	1	5
	Copepoda	1	5		Copepoda	1	5
	Subtotal:	230	1,150		Cyclopoida	1	5
13 Sep 95					Isopoda	1	5
18 m	No. samples processed:	4			Subtotal:	1,176	5,880
	Oligochaeta	396	1,980		Subtotal all depths for this sampling area and date period:	3,317	5,528
	Chironomidae larvae	40	200		Total all dates and depths for Offield:	41,493	3,842
	Ostracoda	3	15				
	Copepoda	2	10				
	Nematoda	1	5				
	Subtotal:	442	2,210				
	Subtotal all depths for this sampling area and date period:	922	1,537				
11 Oct 95							
3 m	No. samples processed:	4					
	Oligochaeta	611	3,055				
	Chironomidae larvae	237	1,185				
	Bivalvia	118	590				
	Nematoda	39	195				
	Ephemeroptera nymph	27	135				
	Prostigmata (water mites)	14	70				
	Cyclopoida	8	40				
	<i>Corophium</i> spp.	4	20				
	Chironomidae pupae	2	10				
	Ostracoda	1	5				
	Turbellaria	1	5				
	Subtotal:	1,062	5,310				
11 Oct 95							
9 m	No. samples processed:	4					
	Oligochaeta	925	4,625				
	Chironomidae larvae	92	460				
	Bivalvia	33	165				

Appendix Table C1. Continued.

Sampling area	Taxon/category	Total number	Density (no./m ²)	Sampling area	Taxon/category	Total number	Density (no./m ²)
Date				Date			
Depth (m)				Depth (m)			
Grand total all sampling areas, dates, depths:		253,045					
Total samples processed (all sampling areas):		647					

Appendix D

Benthic Invertebrates—limnology studies

Processor name: _____

Process Date (dd/mm/yy): _____

Process Time (Hrs.): _____

Collection information

Reservoir: _____

Comments

Date: _____

Site: _____

Replicate no.: _____


ID code: _____

Depth (m): _____


Log ID (from tblWorkLog):_____

[illegible]


Appendix Figure D1. Data analysis sheet used to record organisms recovered from a benthic invertebrate sample.




Appendix Figure D2. Spatial distribution of Oligochaeta on each transect by depth at Silcott Island in Lower Granite Reservoir.




Appendix Figure D3. Spatial distribution of Oligochaeta on each transect by depth at Centennial Island in Lower Granite Reservoir.




Appendix Figure D4. Spatial distribution of Oligochaeta on each transect by depth at Offield in Lower Granite Reservoir.




Appendix Figure D5. Spatial distribution of Chironomidae larvae on each transect by depth at Silcott Island in Lower Granite Reservoir.




Appendix Figure D6. Spatial distribution of Chironomidae larvae on each transect by depth at Centennial Island in Lower Granite Reservoir.




Appendix Figure D7. Spatial distribution of Chironomidae larvae on each transect by depth at Offield in Lower Granite Reservoir.



Appendix Figure D8. Spatial distribution of *Bivalvia* on each transect by depth at Silcott Island in Lower Granite Reservoir.



Appendix Figure D9. Spatial distribution of *Bivalvia* on each transect by depth at Centennial Island in Lower Granite Reservoir.



Appendix Figure D10. Spatial distribution of *Bivalvia* on each transect by depth at Offield in Lower Granite Reservoir.

